## **Supporting Information**

## Fabrication of Conjugated Amphiphilic Triblock Copolymer for Drug Delivery and Fluorescence Cell Imaging

Xuezhi Zhao, Kaicheng Deng, Fangjun Liu, Xiaolong Zhang, Huiru Yang, JinleiPeng, Zengkui Liu, Liwei Ma, Baoyan Wang, Hua Wei\*

State Key Laboratory of Applied Organic Chemistry, Key Laboratory of Nonferrous Metal Chemistry and Resources Utilization of Gansu Province, and College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou, Gansu 730000, China

\*Corresponding author *E-mail address*: weih@lzu.edu.cn (H. Wei)

## List of contents

11 Pages including 1 Scheme and 13 Figures

## **Captions of Scheme and Figure**

Scheme S1. Synthesis of (a) alkyne-POEGMA and (b) PF-b-POEGMA.

Figure S1. SEC elution trace of  $PF_{12}$ -OH using THF as the eluent.

Figure S2.FT-IR spectra of the PF-Br and PF-N<sub>3</sub>.

Figure S3. SEC elution trace of  $PF_{12}$ -OH using DMF as the eluent.

**Figure S4**. SEC elution traces (dRI signals) of alkyne-PCL<sub>44</sub>, alkyne-PCL<sub>44</sub>-b-POEGMA<sub>45</sub>, and PF<sub>12</sub>-b-PCL<sub>44</sub>-b-POEGMA<sub>45</sub>.

Figure S5. <sup>1</sup>H NMR spectrum of alkyne-Br in CDCl<sub>3</sub>.

**Figure S6**.<sup>1</sup>H NMR spectrum of alkyne-POEGMA<sub>45</sub> in CDCl<sub>3</sub>.

Figure S7.<sup>1</sup>H NMR spectrum of PF<sub>12</sub>-*b*-POEGMA<sub>45</sub> in CDCl<sub>3</sub>.

**Figure S8**. Size distributions of (a) P1 and (b) P2 micelles in PBS 7.4 and (c) P1 and (d) P2 micelles in the presence of 10% FBS at a polymer concentration of 0.1 mg/mL determined by DLS.

**Figure S9**. UV–vis absorption and emission spectra of (a) P1, (b) P2 and (c) P0 in water and THF.

Figure S10. Emission spectra of (a) P2 and UV-vis absorption of (b) DOX'HCl in water.

**Figure S11**. UV-vis absorption and emission spectra of (a) DOX@P1, (b) DOX@P2 in water and THF and (c) DOX in water.

**Figure S12**. *In vitro* cytotoxicity of blank P2 micelles in HeLa cells. The blank P2 micelles are almost non-toxic to cells with cell viability above 120% even at the highest tested concentration of 3.8 mg/mL.

Figure S13. In vitro cytotoxicity of free DOX in HeLa cells.



Scheme S1. Synthesis of (a) alkyne-POEGMA and (b) PF-*b*-POEGMA.



Figure S1. SEC elution trace of  $PF_{12}$ -OH using THF as the eluent.



Figure S2. FT-IR spectra of the PF-Br and PF- $N_3$ .



Figure S3. SEC elution trace of  $PF_{12}$ -OH using DMF as the eluent.



**Figure S4**. SEC elution traces (dRI signals) of alkyne-PCL<sub>44</sub>, alkyne-PCL<sub>44</sub>-*b*-POEGMA<sub>45</sub>, and PF<sub>12</sub>-*b*-PCL<sub>44</sub>-*b*-POEGMA<sub>45</sub>.



Figure S5.<sup>1</sup>H NMR spectrum of alkyne-Br in CDCl<sub>3</sub>.



Figure S6.<sup>1</sup>H NMR spectrum of alkyne-POEGMA<sub>45</sub> in CDCl<sub>3</sub>.



**Figure S7**.<sup>1</sup>H NMR spectrum of PF<sub>12</sub>-*b*-POEGMA<sub>45</sub> in CDCl<sub>3</sub>.



**Figure S8**. Size distributions of (a) P1 and (b) P2 micelles in PBS 7.4 and (c) P1 and (d) P2 micelles in the presence of 10% FBS at a polymer concentration of 0.1 mg/mL determined by DLS.



Figure S9. UV-vis absorption and emission spectra of (a) P1, (b) P2 and (c) P0 in



water and THF.

Figure S10. Emission spectra of (a) P2 and UV-vis absorption of (b) DOX'HCl in

water.



Figure S11.UV-vis absorption and emission spectra of (a) DOX@P1, (b) DOX@P2



in water and THF and (c) DOX in water.

**Figure S12**. *In vitro* cytotoxicity of blank P2 micelles in HeLa cells. The blank P2 micelles are almost non-toxic to cells with cell viability above 120% even at the highest tested concentration of 3.8 mg/mL.



Figure S13. In vitro cytotoxicity of free DOX in HeLa cells.