

Synthesis of Heterobimetallic Complexes by Coordination of Rhodium(III) and Iridium(III) Poly-N,O-NHC Complexes to Silver(I), Copper(II) and Zinc(II)

Michael Tegethoff, Florian Roelfes, Christian Schulte to Brinke, Tristan Tsai Yuan Tan, Florian Kampert, Guo-Xin Jin,^b and F. Ekkehardt Hahn^{*}

^aInstitut für Anorganische und Analytische Chemie, Westfälische Wilhelms-Universität Münster, Corrensstraße 28/30, 48149 Münster, Germany

^bDepartment of Chemistry, Fudan University, Shanghai 200433, P. R. China

Table of Contents:

Table of Contents:

1. NMR spectra of [2]	S2
2. NMR spectra of [3]	S3
3. NMR spectra of [4]	S4
4. NMR spectra of [5]	S5
5. NMR spectra of [6]	S6
6. NMR spectra of [10]	S7




Figure S1. ^1H NMR spectrum of [2] in CDCl_3 .




Figure S2. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of [2] in CDCl_3 .




Figure S3. ^1H NMR spectrum of [3] in CDCl_3 .




Figure S4. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of [3] in CDCl_3 .




Figure S5. ^1H NMR spectrum of [4] in CDCl_3 .




Figure S6. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of [4] in CDCl_3 .




Figure S7. ^1H NMR spectrum of [5] in CD_2Cl_2 .




Figure S8. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of [5] in CD_2Cl_2 . The resonance for the carbene carbon atoms (C2) was not observed.




Figure S9. ^1H NMR spectrum of [6] in CD_3OD .




Figure S10. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of [6] in CD_3OD .




Figure S11. ^1H NMR spectrum of [10] in CDCl_3 .




Figure S12. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of [10] in CDCl_3 .