
Supporting Information


A domino N₂-extrusion-cyclization of alkynylarylketone derivatives for the synthesis of indoloquinolines and carbocyclefused quinolines


Bhornrawin Akkachairin,[†] Jumreang Tummatorn,* ^{†‡} Narumol Khamsuwan[‡] Charnsak Thongsornkleeb^{†‡} and Somsak Ruchirawat^{†‡}


 [†]Program on Chemical Biology, Chulabhorn Graduate Institute, Center of Excellence on Environmental Health and Toxicology (EHT), Ministry of Education, 54 Kamphaeng Phet 6, Laksi, Bangkok 10210, Thailand
[‡]Laboratory of Medicinal Chemistry, Chulabhorn Research Institute, 54 Kamphaeng Phet 6, Laksi, Bangkok 10210, Thailand


	Contents:	Pages
1.	Copies of ¹ H and ¹³ C- NMR spectra for compound 1a-1x	S2-S49
2.	Copies of ¹ H and ¹³ C- NMR spectra for compound 2a-2x' and 3a	S50-S111
3.	Copies of ¹ H and ¹³ C- NMR spectra for compound 8a-8q and 9a-9b	S112-S151
4.	Copies of ¹ H and ¹³ C- NMR spectra for compound 4a-4r and 6a-6b	S152-S191
5.	Copies of ¹ H and ¹³ C- NMR spectra for compound 5a-5r and 7a-7b	S192-S239
6.	Optimazation of Method for the synthesis of indoloquinolines (Table S1)	S240
7.	Optimazation of Method for the synthesis of carbocycle-fused quinolines (Table S2)	S241






















































































































































- 14.06













S54








S56


















S62








-18.93
























































S82







45.78

12.66 49

d


4 5


61


42.19


42.17


36.














































































































































































































































































































































































































































































Optimization of Method for the synthesis of indoloquinolines

Ortho-alkynylarylketone **1a** was employed as the substrate to screen for the optimal condition. To determine the ratio between indoloquinoline product and 1, 5-diketone by-product, we varied equivalent of TMSN₃ and TfOH as well as solvent and concentration to improve the yield of the desired product and reduce the formation of by-product. The isolated yield of compound **2a** and **3a** in each entry is summarized in the table below.




Table S1. Optimization for the synthesis of indoloquinoline 2a from alkynylarylketone 1a.*a*Isolated yields.

Entry	Compound 1a (equiv)	TMSN3 (equiv)	TfOH (equiv)	Solvent	%yield ^a	
				(dry)	2a	3 a
1	1.0	2.5	2.5	0.1 M DCM	48	22
2	1.0	3.0	3.0	0.1 M DCM	47	16
3	1.0	4.0	4.0	0.1 M DCM	50	19
4	1.0	3.5	3.5	0.1 M DCM	49	23
5	1.0	3.5	3.5	0.2 M DCM	54	13
6	1.0	3.5	3.5	0.2 M Toluene	46	14
7	1.0	3.5	3.5	0.3 M DCM	58	8

Optimization of Method for the synthesis of carbocycle-fused quinolines

To screen for the optimal condition, α -alkynylarylketone **4a** was used as the precursor. Compound **4a** was treated under the optimal condition of indoloquinoline method, resulting moderate yield of desired product **5a**. In order to improve the yield of product, we varied various variables (TMSN₃, TfOH, solvents and concentrations, time and temperature. The result of the optimizations is shown in Table 2.



Table S2. Optimization for the carbocycle-fused quinoline **5a** from α -alkynylarylketone **4a**. ^{*a*}Isolated yields.

Entry	Compound 4a (equiv)	TMSN ₃ (equiv)	TfOH (equiv)	Solvent (dry)	Time	%yield ^a
1	1.0	3.5	3.5	0.3 M DCM	O/N, rt	58
2	1.0	1.2	1.2	0.1 M DCM	1 h, rt	77
2	1.0	1.2	1.1	0.1 M Tolulene	1 h, rt	54
3	1.0	1.2	1.2	0.05 M DCM	1 h, rt	71
4	1.0	1.2	1.2	0.2 M DCM	1 h, rt	75
5	1.0	1.2	1.1	0.1 M DCM	1 h, rt	65
6	1.0	1.0	1.0	0.1 M DCM	1 h, rt	63
7	1.0	1.2	1.2	0.1 M DCM	3 h, rt	72
8	1.0	1.4	1.4	0.1 M DCM	1 h, rt	75
9	1.0	2.0	2.0	0.1 M DCM	1 h, 0 °C	67