

Supporting Information

Determination and Correlation of the Solubility of L-Fucose in Four Binary Solvent Systems at the Temperature Range from 288.15 K to 308.15 K

Jingxuan Qiu[†], Shuang Song[†], Xinglong Chen[†], Dengjing Yi^{†§}, Mengyao An^{‡§}, and Peng Wang^{*†§}

[†] School of Chemical Engineering, [‡] School of Chemical and Life Science, and [§] Advanced Institute of Materials Science Changchun University of Technology, Changchun, Jilin 130012, People's Republic of China

Corresponding Author

*Tel: +86-431-85968101, Fax: +86-431-85968101, Email: pengwang@ccut.edu.cn

Figure Captions

Figure S1. Experimental (x_A^{exp}) solubility scatter diagram of L -fucose in methanol + water at the temperature range from 288.15 K to 308.15 K

Figure S2. Experimental (x_A^{exp}) solubility scatter diagram of L -fucose in ethanol + water at the temperature range from 288.15 K to 308.15 K

Figure S3. Experimental (x_A) solubility scatter diagram of L -fucose in 2-propanol + water at the temperature range from 288.15 K to 308.15 K

Figure S4. Experimental (x_A) solubility scatter diagram of L -fucose in acetone + water at the temperature range from 288.15 K to 308.15 K

Figure S5. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for L -fucose solubility in the full range in methanol + water

Figure S6. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for L-fucose solubility in the full range in ethanol + water

Figure S7. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for L-fucose solubility in the full range in 2-propanol + water

Figure S8. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for L-fucose solubility in the full range in acetone + water




Figure S1. Experimental (x_A^{exp}) solubility scatter diagram of $_L$ -fucose in methanol + water at the temperature range from 288.15 K to 308.15 K




Figure S2. Experimental (x_A^{exp}) solubility scatter diagram of $_L$ -fucose in ethanol + water at the temperature range from 288.15 K to 308.15 K




Figure S3. Experimental (x_A) solubility scatter diagram of L -fucose in 2-propanol + water at the temperature range from 288.15 K to 308.15 K




Figure S4. Experimental (x_A) solubility scatter diagram of L -fucose in acetone + water at the temperature range from 288.15 K to 308.15 K




Figure S5. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for $_L$ -fucose solubility in the full range in methanol + water




Figure S6. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for $_L$ -fucose solubility in the full range in ethanol + water




Figure S7. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for L -fucose solubility in the full range in 2-propanol + water




Figure S8. Apelblat-Jouyban-Acree model fitting plot of $\ln(x_A)$ versus T (from 288.15 K to 308.15 K) and x_1 (from 0.0 to 1.0) for L -fucose solubility in the full range in acetone + water