Supporting Information

Fluorescent ZnS quantum dots-phosphoethanolamine nanoconjugates for bioimaging live cells in cancer research

Alexandra A. P. Mansur[a], Herman S. Mansur[a,*], Sandhra M. Carvalho[a,b,c], Zélia I. P. Lobato[b], Maria de Fátima Leite[c], Lorena L. Mansur[a,b]

[a]*Center of Nanoscience, Nanotechnology and Innovation - CeNano²I, Federal University of Minas Gerais-UFMG, Av. Antônio Carlos, 6627 – Belo Horizonte/MG, Brazil.*
e-mail: hmansur@demet.ufmg.br (HSM)
[b]*Department of Preventive Veterinary Medicine, Veterinary School, Federal University of Minas Gerais-UFMG, Av. Antônio Carlos, 6627 – Belo Horizonte/MG, Brazil.*
e-mail: ziplobato@gmail.com (ZIPL)
[c]*Department of Physiology and Biophysics, ICB, Federal University of Minas Gerais-UFMG, Av. Antônio Carlos, 6627 – Belo Horizonte/MG,Brazil.*
e-mail: leitemd@ufmg.br (MFL)



Figure S1 – Cell viability responses by MTT assay after 48 h of incubation with the PEA and DOX at concentration of 0.1 μM.

* To whom correspondence should be addressed: Federal University of Minas Gerais, Av. Antônio Carlos, 6627 – Escola de Engenharia, Bloco 2 – Sala 2233, 31.270-901, Belo Horizonte/MG, Brazil; TelFax: +55-31-34091843; E-mail: hmansur@demet.ufmg.br (H. Mansur)



Figure S2 – Cell viability responses by MTT assay after 48 h of incubation with the PEA and DOX at concentration of 1.0 μM.



Figure S3 – Cell viability responses by MTT assay after 48 h of incubation with the PEA and DOX at concentration of 10 μM.



Figure S4 – Cell viability responses by MTT assay after 48 h of incubation with the PEA and DOX at concentration of 100 μM.