

Supporting Information

Radiolysis Induced Formation of Hollow RbBr Nanocubes under Electron Beam Irradiation

Linyuan Lian¹, Feng Cheng², Yong Xia¹, Moyan Zheng¹, Jiangping Ke¹, Weizhuo Zhang¹, Huan Liu¹, Daoli Zhang¹, Jianbo Gao³, Jiang Tang², Luying Li², Manlin Tan^{4} and Jianbing Zhang^{1,5*}*

¹School of Optical and Electronic Information, Engineering Research Center for Functional Ceramics, Ministry of Education, Huazhong University of Science and Technology, 1037 Luoyu Road, Wuhan, Hubei 430074, China

²Wuhan National Laboratory for Optoelectronics, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China

³Department of Physics and Astronomy, Clemson University, South Carolina 29634, United States

⁴Research Institute of Tsinghua University in Shenzhen, Shenzhen, Guangdong 518057, China

⁵Shenzhen Huazhong University of Science and Technology Research Institute, Shenzhen, Guangdong 518057, China




Figure S1. The energy-dispersive X-ray spectroscopy result of the hollow RbBr nanocubes.




Figure S2. The HAADF image of the RbBr nanocubes showing voids only appeared in the area with focused e-beam.