Spectral, Electrochemical and ESR Characterization of Manganese Tetraarylporphyrins Containing Four β,β'-Pyrrole Fused Butano and Benzo Groups in Nonaqueous Media

Yuanyuan Fang*[a], Liping Wang[a], Weijie Xu[a], Zhongping Ou[b], Mingyuan Chen[a,b], Lei Cong[b], Wenqian Shan[b], Xiangyi Ke[b] and Karl M. Kadish*[b]

[a]*School of Chemistry and Chemical Engineering, Jiangsu University, Zhenjiang, 212013, China*
[b]*Department of Chemistry, University of Houston, Houston, TX 77204-5003, USA*

Content	Page
Figure S1. UV-vis spectra of $(Ar)_4$PorMnCl **1a** and **1b** in (a) CH_2Cl_2 and (b) pyridine, 0.1 M TBAP.	2
Figure S2. UV-vis spectra of (a) butano$(Ar)_4$PorMnCl **2a**, **2b** and (b) benzo$(Ar)_4$PorMnCl **3a, 3b** in CH_2Cl_2, 0.1 M TBAP.	3
Figure S3. UV-vis spectra of (a) butano$(Ar)_4$PorMnCl **2a**, **2b** and (b) benzo$(Ar)_4$PorMnCl **3a**, **3b** in pyridine, 0.1 M TBAP.	4
Figure S4. Cyclic voltammograms showing the reductions of (a) $(Ar)_4$PorMnCl **1a**, **1c**, (b) butano$(Ar)_4$PorMnCl **2a**, **2c** and (c) benzo$(Ar)_4$PorMnCl **3a**, **3c** (Ar = *p*-CH_3Ph for **1a-3a**, Ar = *p*-ClPh for **1c-3c**) in CH_2Cl_2, 0.1 M TBAP at a scan rate of 0.10 V/s.	5
Figure S5. Cycic voltammograms showing oxidations of (a) butano$(Ar)_4$PorMnCl **2a-2c** and (b) benzoo$(Ar)_4$PorMnCl **3a-3c** (Ar = *p*-CH_3Ph for **2a**, **3a**, Ar = Ph for **2b, 3b** and Ar = *p*-ClPh for **2c**, **3c**) in CH_2Cl_2, 0.1 M TBAP at a scan rate of 0.10 V/s.	6
Figure S6. Cyclic voltammograms showing the reductions of (a) $(Ar)_4$PorMnCl **1a**, **1c** and (b) butano$(Ar)_4$PorMnCl **2a**, **2c** (Ar = *p*-CH_3Ph for **1a** and **2a**, Ar = *p*-ClPh for **1c** and **2c**) in pyridine, 0.1 M TBAP at a scan rate of 0.10 V/s.	7
Figure S7. Cyclic voltammograms showing the reductions of (a) benzo$(CH_3Ph)_4$PorMnCl **3a** and (b) benzo$(ClPh)_4$PorMnCl **3c** in pyridine, 0.1 M TBAP at a scan rate of 0.10 V/s.	8
Figure S8. UV-vis spectral changes of (a) butano$(CH_3Ph)_4$PorMnCl **2a** and (b) butano$(Ph)_4$PorMnCl **2b** during controlled potential reductions in pyridine containing 0.1 M TBAP.	9



Figure S1. UV-visible spectra of $(Ar)_4$PorMnCl **1a** and **1b** in (a) CH_2Cl_2 and (b) pyridine, 0.1 M TBAP.



Figure S2. UV-vis spectra of (a) butano(Ar)$_4$PorMnCl **2a** and **2b** and (b) benzo(Ar)$_4$PorMnCl **3a** and **3b** in CH_2Cl_2, 0.1 M TBAP.



Figure S3. UV-vis spectra of (a) butano(Ar)$_4$PorMnCl **2a** and **2b** and (b) benzo(Ar)$_4$PorMnCl **3a** and **3b** in pyridine, 0.1 M TBAP.



Figure S4. Cyclic voltammograms showing the reductions of (a) $(Ar)_4$PorMnCl **1a**, **1c**, (b) butano$(Ar)_4$PorMnCl **2a**, **2c** and (c) benzo$(Ar)_4$PorMnCl **3a**, **3c** (Ar = CH_3Ph for **1a**-**3a**, Ar = ClPh for **1c**-**3c**)in CH_2Cl_2, 0.1 M TBAP at a scan rate of 0.10 V/s.



Figure S5. Cycic voltammograms showing oxidations of (a) butano(Ar)$_4$PorMnCl **2a-2c** and (b) benzoo(Ar)$_4$PorMnCl **3a-3c** (Ar = CH_3Ph for **2a, 3a**, Ar = Ph for **2b, 3b** and Ar = ClPh for **2c, 3c**) in CH_2Cl_2, 0.1 M TBAP at a scan rate of 0.10 V/s.



Figure S6. Cyclic voltammograms showing the reductions of (a) $(Ar)_4$PorMnCl **1a**, **1c** and (b) butano$(Ar)_4$PorMnCl **2a**, **2c** (Ar = CH_3Ph for **1a** and **2a**, Ar = ClPh for **1c** and **2c**) in pyridine, 0.1 M TBAP at a scan rate of 0.10 V/s.



Figure S7. Cyclic voltammograms showing the reductions of (a) benzo(CH_3Ph)$_4$PorMnCl **3a** and (b) benzo(ClPh)$_4$PorMnCl **3c** in pyridine, 0.1 M TBAP at a scan rate of 0.10 V/s.



Figure S8. UV-vis spectral changes of (a) butano($CH_3Ph)_4$PorMnCl **2a** and (b) butano$(Ph)_4$PorMnCl **2b** during controlled potential reductions in pyridine containing 0.1 M TBAP.