

Confinement and Time Immemorial: Prebiotic Synthesis of Nucleotides on a Porous Mineral Nanoreactor

Francisco Rodrigues,^{a,b,*} Thomas Georgelin,^c Guillaume Gabant,^c Baptiste Rigaud,^d Fabrice Gaslain,^e

Guanzheng Zhuang,^a Maria Gardênnia da Fonseca,^f Valentin Valtchev^g, David Touboul^h and Maguy Jaber^{a,*}

^a Sorbonne University, CNRS UMR 8220, Laboratoire d'Archéologie Moléculaire et Structurale, 4 place Jussieu, F-75005 Paris, France.

^b Department of Chemistry, State University of Paraíba, UEPB, Campina Grande, Paraíba, Brazil.

^c Centre de Biophysique Moléculaire, CNRS, Rue Charles Sadron, 45000 Orléans, France.

^d CNRS Institut des Matériaux de Paris Centre (FR2482), 4 place jussieu, 75005 Paris, France.

^e MINES ParisTech, PSL - Research University, MAT - Centre des matériaux, CNRS UMR 7633, BP 87, F-91003 Evry, France.

^f Department of Chemistry, Federal University of Paraíba, UFPB, João Pessoa, Paraíba, Brazil.

^g Normandy University, Laboratoire Catalyse & Spectrochimie, ENSICAEN, 6 bl Maréchal Juin, 14050 Caen, France.

^h CNRS-Institut de Chimie des Substances Naturelles, UPR2301, Université Paris-Saclay, 91198 Gif-sur-Yvette cedex, France


AUTHOR INFORMATION

Corresponding Author


*francisco.rodrigues@cct.uepb.edu.br ; maguy.jaber@upmc.fr

Supporting Information - SI


SI-1. N₂ adsorption and desorption isotherms of (a) parent Na-Mordenite and (b) (R+Pi)A@MOR.




SI-2. XRD pattern for the bulk mordenite (MOR) and (R+P)A@MOR samples before and after activation.



SI-3. ^{27}Al NMR MAS spectra for the bulk mordenite (MOR) and (R+P)A@MOR samples before and after activation.



SI-4. ^{13}C CP NMR spectra of bulk ribose (R).



*f and p stand for furanose and pyranose, respectively.