Supporting Information

Antitumor Activity of Vanicoside B Isolated from *Persicaria dissitiflora* by Targeting CDK8 in Triple-Negative Breast Cancer Cells

Donghwa Kim,[†] Cai Yi Wang,[†] Ruoci Hu,[†] Ji Yun Lee,[†] Thi-Thu-Trang Luu,[†] Hee-Juhn Park,^{*,‡} and Sang Kook Lee^{*,†}

[†]College of Pharmacy, Natural Products Research Institute, Seoul National University, Seoul 08826, Republic of Korea
[‡]Department of Pharmaceutical Engineering, Sangji University, Wonju 26339, Republic of Korea

Contents

Table S1. Inhibitory effects of 1–3 on the proliferation of five human cancer cell lines......1

IC ₅₀ (µM) ^a			Cell line ^c		
compound	SK-Hep-1	A549	HCT116	SNU638	MDA-MB-231
quercitrin (1)	>10	>10	>10	>10	>10
vanicoside B (2)	>10	>10	>10	>10	9.0
lapathoside C (3)	>10	>10	>10	>10	>10
etoposide ^b	0.39	0.15	0.89	0.55	4.0

Table S1. Inhibitory Effects of 1–3 on the Proliferation of Five Human Cancer Cell Lines

^{*a*}Results were expressed as the calculated half maximal inhibitory concentration of **1-3** and etoposide (μ M). ^{*b*}Etoposide was used as a positive control. ^{*c*}Human cancer cell lines were SK-Hep1 (human liver cancer cell), A549 (human lung cancer cell), HCT116 (human colon cancer cell), SNU638 (human stomach cancer cell), and MDA-MB-231 (human breast cancer cell).