Supporting Information

Organocatalytic Asymmetric Hetero-Diels–Alder Reaction of in-situ Generated Dienes: Access to α,β -Unsaturated δ -Lactones Featuring CF₃-Substituted Quaternary Stereocenter

Harshit Joshi, Ankit Yadav, Arko Das, and Vinod K. Singh*


Department of Chemistry, Indian Institute of Technology Kanpur, Kanpur, UP – 208 016, India

Email: vinodks@iitk.ac.in

Table of contents:

1.	X-ray crystal structure and crystallographic data of 4ac	S2-S3
2.	X-ray crystal structure and crystallographic data of 6	S4-S5
3.	References	S5
4.	NMR spectra	S6-S51
5.	HPLC graph	S52-S71

1. X-ray crystal structure and crystallographic data of 4ac (CCDC 1951698):



Sample preparation: 50.0 mg of compound **4ac** was taken in a vial and dissolved in 5 mL of ethyl acetate then 0.2 mL n-hexane was added to it (dropwise). The solvents were allowed to evaporate slowly at room temperature, which yielded block-shaped transparent crystals of compound **4ac** in two weeks.




Figure S1: Molecular structure of **4ac** with ellipsoids set at 50% probability. All hydrogen atoms except N-H are omitted for clarity.

Crystal structure determination for compound 4ac: The crystal structure was solved by SHELXT^{1a} structure solution program and refined by full matrix least-squares method based on F^2 by SHELXL^{1b} program in the Olex 2^{1c} software. All the crystallographic figures were generated using the Diamond 3.2k software.

Table S1: Crystal data for **4ac**

Identification code	4ac		
Empirical formula	C ₂₅ H ₁₇ BrF ₃ NO ₃		
Formula weight	516.30		
Temperature/K	273.15		
Crystal system	monoclinic		
Space group	P2 ₁		
a/Å	9.8347(5)		
b/Å	22.0058(12)		
c/Å	10.1652(6)		
α/°	90		
β/°	94.998(2)		
γ/°	90		
Volume/Å ³	2191.6(2)		
Z	4		
$\rho_{calc} g/cm^3$	1.565		
μ /mm ⁻¹	1.929		
F(000)	1040.0		
Crystal size/mm ³	$0.052 \times 0.023 \times 0.015$		
Radiation	MoKα ($\lambda = 0.71073$)		
2Θ range for data collection/° 5.468 to 56.556			
Index ranges	$-13 \le h \le 13$, $-29 \le k \le 29$, $-13 \le l \le 13$		
Reflections collected	33659		
Independent reflections	10757 [$R_{int} = 0.0466, R_{sigma} = 0.0610$]		
Data/restraints/parameters	10757/1/595		
Goodness-of-fit on F ²	1.013		
Final R indexes [I>= 2σ (I)]	$R_1 = 0.0293, wR_2 = 0.0685$		
Final R indexes [all data]	$R_1 = 0.0329, wR_2 = 0.0701$		
Largest diff. peak/hole / e Å-3 0.86/-0.58			
Flack parameter	0.034(3)		

2. X-ray crystal structure and crystallographic data of 6 (CCDC 1951699):

Ph. OH NHBz
F₃C (R)

6

Sample preparation: 30.0 mg of compound **6** was taken in a vial and dissolved in 5 mL of ethyl acetate then 0.2 mL n-hexane was added to it in a dropwise manner. The solvents were allowed to evaporate slowly at room temperature, which yielded block-shaped transparent crystals of compound **6** in one week.




Figure S2: Molecular structure of **6** with ellipsoids set at 50% probability. All hydrogen atoms except N-H and O-H are omitted for clarity.


Crystal structure determination for compound 6: The crystal structure was solved by SHELXT^{1a} structure solution program and refined by full matrix least-squares method based on F^2 by SHELXL^{1b} program in the Olex 2^{1c} software. All the crystallographic figures were generated using the Diamond $3.2k^{1d}$ software.


Table S2: Crystal data for compound 6

Identification code	6		
Empirical formula	$C_{29}H_{26}BrF_3N_2O_3$		
Formula weight	587.42		
Temperature/K	273.15		
Crystal system	orthorhombic		
Space group	P2 ₁ 2 ₁ 2 ₁		
a/Å	10.6343(5)		
b/Å	11.7382(5)		
c/Å	20.6645(10)		
α/°	90		
β/°	90		
γ/°	90		
Volume/Å ³	2579.5(2)		
Z	4		
$\rho_{calc}g/cm^3$	1.513		
μ /mm ⁻¹	1.650		
F(000)	1200.0		
Crystal size/mm ³	$0.029 \times 0.026 \times 0.023$		
Radiation	ΜοΚα (λ = 0.71073)		
2Θ range for data collection/° 5.532 to 51.988			
Index ranges	$-13 \le h \le 13$, $-14 \le k \le 14$, $-25 \le l \le 25$		
Reflections collected	34484		
Independent reflections	$5062 [R_{int} = 0.0841, R_{sigma} = 0.0541]$		
Data/restraints/parameters	5062/0/344		
Goodness-of-fit on F ²	1.064		
Final R indexes [I>= 2σ (I)]	$R_1 = 0.0292, wR_2 = 0.0636$		
Final R indexes [all data]	$R_1 = 0.0375, wR_2 = 0.0688$		
Largest diff. peak/hole / e Å-3 0.26/-0.30			
Flack parameter	0.022(5)		


3. References: 1 (a) Sheldrick, G. M. SHELXT - Integrated Space-Group and Crystal-Structure Determination. *Acta Cryst.* 2015, *A71*, 3. (b) Sheldrick, G. M. Crystal Structure Refinement with SHELXL. *Acta Cryst.* 2015, *C71*, 3. (c) Dolomanov, O. V.; Bourhis, L. J.; Gildea, R. J.; Howard, J. A. K.; Puschmann, H. OLEX2: A Complete Structure Solution, Refinement and Analysis Program. *J. Appl. Crystallogr.* 2009, *42*, 339. (d) Brandenburg, K.; DIAMOND. Version 3.2k, Crystal Impact GbR, Bonn, Germany, 2014.

4. NMR specta:







 $400~\text{MHz}\ ^1\text{H}\ \text{NMR}$ and $100~\text{MHz}\ ^{13}\text{C}\ \text{NMR}$ spectra of compound 3d in $CDCl_3$



400 MHz 1 H NMR and 100 MHz 13 C NMR spectra of compound 3k, $\it E/Z$ (1:1.3), in CDCl $_3$




400 MHz ¹H NMR and 100 MHz ¹³C NMR spectra of compound **3m**, E/Z (1:1.1), in CDCl₃




400 MHz ^1H NMR and 100 MHz ^{13}C NMR spectra of compound 3n, E/Z (1:1.3), in CDCl $_3$

NMR spectra of HDA adducts (4), transformation products (5) and (6)

(i) $(R)-N-(2-\infty -4,6-\text{diphenyl-}6-(\text{trifluoromethyl})-5,6-\text{dihydro-}2H-\text{pyran-}3-\text{yl})$ benzamide (4aa)




500 MHz ¹H NMR spectrum of compound **4aa** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4aa in $CDCl_3$

$\hbox{$(ii)$} \qquad \hbox{$(R)$-$N$-$(4-(3-bromophenyl)-2-oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) benzamide $(4ab)$ }$




500 MHz ¹H NMR spectrum of compound **4ab** in CDCl₃




125 MHz ^{13}C NMR and 373 MHz ^{19}F NMR spectra of compound 4ab in CDCl $_3$


(iii) (R)-N-(4-(4-bromophenyl)-2-oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) benzamide (4ac)




500 MHz ¹H NMR spectrum of compound **4ac** in CDCl₃



 $125\,MHz$ ^{13}C NMR and 373 MHz ^{19}F NMR spectra of compound $\boldsymbol{4ac}$ in $CDCl_3$




 $500\ MHz\ ^1H\ NMR\ spectrum\ of\ compound\ \mbox{\bf 4ad}$ in $CDCl_3$




 $125\,MHz$ ^{13}C NMR and 373 MHz ^{19}F NMR spectra of compound $\boldsymbol{4ad}$ in $CDCl_3$

$(v) \qquad (R)-N-(4-(4-\text{chlorophenyl})-2-\text{oxo-}6-\text{phenyl-}6-(\text{trifluoromethyl})-5,6-\text{dihydro-}2H-\text{pyran-}3-\text{yl})\\ \text{benzamide } (4\text{ae})$




500 MHz ^1H NMR spectrum of compound 4ae in CDCl_3




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4ae in $CDCl_3$

$(vi) \qquad (R)-N-(4-(4-fluorophenyl)-2-oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) \\ benzamide~(4af)$




500 MHz 1 H NMR spectrum of compound ${\bf 4af}$ in CDCl $_3$




125 MHz ¹³C NMR and 373 MHz ¹⁹F NMR spectra of compound **4af** in CDCl₃

$(vii) \qquad (R)-N-(4-(4-nitrophenyl)-2-oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) \\ benzamide~(4ag)$




500 MHz ¹H NMR spectrum of compound **4ag** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4ag in $CDCl_3$

$(viii) \qquad (R)-N-(2-oxo-6-phenyl-4-(p-tolyl)-6-(trifluoromethyl)-5, 6-dihydro-2H-pyran-3-yl) benzamide \ (4ah)$




500 MHz ¹H NMR spectrum of compound **4ah** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound $\boldsymbol{4ah}$ in $CDCl_3$

$\hbox{$(ix)$} \qquad \hbox{$(R)$-$N-$(4-(4-methoxyphenyl)-2-oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2$H-pyran-3-yl)} benzamide \hbox{$(4ai)$}$




500 MHz ¹H NMR spectrum of compound **4ai** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4ai in $CDCl_3$

$(x) \qquad (R)-N-(2-oxo-6-phenyl-4-(thiophen-2-yl)-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) \\ benzamide~(4aj)$




500 MHz ¹H NMR spectrum of compound **4aj** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4aj in $CDCl_3$

$(xi) \qquad (R)-N-(4-(benzofuran-2-yl)-2-oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) \\ benzamide~(4ak)$





500 MHz 1 H NMR spectrum of compound $\mathbf{4ak}$ in CDCl $_3$



 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4ak in $CDCl_3$


$(\textbf{xii}) \qquad (\textbf{\textit{R}}) - \textbf{\textit{N}} - (\textbf{\textit{4}} - (\textbf{\textit{naphthalen-2-yl}}) - \textbf{\textit{2}} - \textbf{\textit{oxo-6-phenyl-6-(trifluoromethyl)-5,6-dihydro-2} H - \textbf{\textit{pyran-3-yl}}) \\ \text{benzamide (4al)}$






100 MHz 13 C NMR and 373 MHz 19 F NMR spectra of compound **4al** in CDCl₃

$\begin{array}{ll} \hbox{(xiii)} & \hbox{(R)-$N-$(4-cyclopropyl-2-oxo-6-phenyl-$6-$(trifluoromethyl)-$5,6-dihydro-$2$H-pyran-$3$-yl)} \\ & \hbox{benzamide (4am)} \end{array}$




500 MHz ¹H NMR spectrum of compound **4am** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound $\boldsymbol{4am}$ in $CDCl_3$

$\begin{array}{ll} \hbox{(xiv)} & \hbox{(R)-$N-$(4-isopropyl-2-oxo-6-phenyl-6-(trifluoromethyl)-$5,6$-dihydro-2H-pyran-3-yl)} \\ & \hbox{benzamide (4an)} \end{array}$




500 MHz ¹H NMR spectrum of compound **4an** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound $\boldsymbol{4an}$ in $CDCl_3$

$(xv) \qquad (R)-N-(6-(4-{\rm fluorophenyl})-2-{\rm oxo}-4-{\rm phenyl}-6-({\rm trifluoromethyl})-5, \\ 6-{\rm dihydro}-2H-{\rm pyran}-3-{\rm yl}) \\ {\rm benzamide}~(4{\rm ba})$




500 MHz ¹H NMR spectrum of compound **4ba** in CDCl₃




 $100\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound $\mbox{\bf 4ba}$ in $CDCl_3$

$(xvi) \qquad (R)-N-(6-(4-chlorophenyl)-2-oxo-4-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) \\ benzamide~(4ca)$




500 MHz ¹H NMR spectrum of compound **4ca** in CDCl₃




 $125\,MHz$ ^{13}C NMR and 373 MHz ^{19}F NMR spectra of compound 4ca in $CDCl_3$

(xvii) (R)-N-(2-oxo-4-phenyl-6-(p-tolyl)-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) benzamide (4da)




500 MHz ¹H NMR spectrum of compound **4da** in CDCl₃




 $125\,MHz$ ^{13}C NMR and 373 MHz ^{19}F NMR spectra of compound 4da in $CDCl_3$

 $(xviii) \quad (R)-N-(6-(2,3-dihydrobenzofuran-5-yl)-2-oxo-4-phenyl-6-(trifluoromethyl)-5,6-dihydro-2H-pyran-3-yl) benzamide (4ea)$




500 MHz ¹H NMR spectrum of compound **4ea** in CDCl₃




 $125\ MHz\ ^{13}C\ NMR$ and $373\ MHz\ ^{19}F\ NMR$ spectra of compound 4ea in $CDCl_3$

$(xix) \qquad (R)-N-(6-(4-\text{chlorophenyl})-4-(4-\text{methoxyphenyl})-2-\text{oxo-}6-(\text{trifluoromethyl})-5,6-\text{dihydro-}2H-\\ \text{pyran-}3-\text{yl})\text{benzamide } (4\text{ci})$




500 MHz ¹H NMR spectrum of compound **4ci** in CDCl₃




125 MHz 13 C NMR and 373 MHz 19 F NMR spectra of compound 4ci in CDCl $_3$

$(xx) \qquad (R,Z)-N-(3-(4-bromophenyl)-6,6,6-trifluoro-1-hydrazinyl-5-hydroxy-1-oxo-5-phenylhex-2-en-2-yl) benzamide (5)$




400 MHz ¹H NMR spectrum of compound **5** in DMSO-*d6*




100 MHz $^{13}\mathrm{C}$ NMR and 373 MHz $^{19}\mathrm{F}$ NMR spectra of compound 5 in DMSO-d6

$(xxi) \qquad (\textit{R,Z})-\textit{N-}(3-(4-bromophenyl)-6,6,6-trifluoro-5-hydroxy-1-oxo-5-phenyl-1-(pyrrolidin-1-yl)hex-2-en-2-yl)benzamide (6)$




500 MHz ¹H NMR spectrum of compound **6** in DMSO-*d6*




100 MHz 13 C NMR and 373 MHz 19 F NMR spectra of compound **6** in DMSO-d6

5. HPLC graphs:




				Area [mAU*s]	Height [mAU]	
1	35.169	MM	2.8966	9621.98242	55.36385	50.7781
2	53.191	MM	4.3123	9327.10547	36.04860	49.2219

HPLC graph of racemic 4aa




Peak :	RetTime	Type	Width	Area	Height	Area	
#	[min]		[min]	[mAU*s]	[mAU]	%	
1	35.612	MM	2.5148	2184.18701	14.47580	1.5735	
2	52.308	MM	4.1496	1.36627e5	548.75946	98.4265	

HPLC graph of enantioenriched 4aa



Peak	RetTime	Туре	Width	Area	Height	Area	
#	[min]		[min]	[mAU*s]	[mAU]	8	
1	24.961	MM	1.9580	4.39370e4	373.99213	50.4353	
2	34.487	MM	2.7986	4.31786e4	257.14069	49.5647	

HPLC graph of racemic 4ab




HPLC graph of enantioenriched **4ab**

136.01842


98.8869

2.8899 2.35850e4


34.605 MM



HPLC graph of racemic 4ac




HPLC graph of enantioenriched 4ac




[min] [min] [mAU*s] [mAU] %
----|-----|-----|------|
1 24.297 MF 1.9373 4.27175e4 367.50412 50.1487
2 33.696 FM 2.7735 4.24642e4 255.17612 49.8513


HPLC graph of racemic 4ad




HPLC graph of enantioenriched 4ad




HPLC graph of racemic 4ae



HPLC graph of enantioenriched 4ae



HPLC graph of racemic 4af




HPLC graph of enantioenriched 4af

3.9295 6.49443e4


275.45367

2 52.891 MM




Peak	RetTime	Type	Width	Area	Height	Area	
#	[min]		[min]	[mAU*s]	[mAU]	8	
1	61.988	MF	4.8131	7.55600e4	261.64972	48.4634	
2	79.641	FM	6.0872	8.03517e4	220.00171	51.5366	

HPLC graph of racemic 4ag




HPLC graph of enantioenriched 4ag




1 39.021 MF 3.0157 1.62827e4 89.98800 49.7682 2 53.288 FM 4.4473 1.64344e4 61.58958 50.2318


HPLC graph of racemic 4ah




HPLC graph of enantioenriched 4ah




HPLC graph of racemic 4ai




HPLC graph of enantioenriched 4ai



HPLC graph of racemic 4aj




HPLC graph of enantioenriched 4aj



	RetTime			Area [mAU*s]	Height [mAU]	Area %
1	12.733	MM	0.4914	1.30234e4	441.73514	50.2709
2	21.188	MM	0.8203	1.28830e4	261.74518	49.7291

HPLC graph of racemic 4ak



HPLC graph of enantioenriched 4ak

0.8451 9.82552e4

0.5095


29.90906

1937.71313


0.9219

1 12.806 MM


2 21.281 MM



HPLC graph of racemic 4al




HPLC graph of enantioenriched 4al




Peak	RetTime	Type	Width	Area	Height	Area
#	[min]		[min]	[mAU*s]	[mAU]	%
1	20.866	MF	0.6547	5652.54248	143.89999	49.6984
2	23.042	FM	0.7745	5721.14453	123.11797	50.3016

HPLC graph of racemic 4am




HPLC graph of enantioenriched 4am




Peak	RetTime	Type	Width	Area	Height	Area	
#	[min]		[min]	[mAU*s]	[mAU]	용	
1	15.152	MF	1.1545	5242.55908	75.68300	47.4055	
2	17.900	FM	1.4935	5816.40527	64.90762	52.5945	

HPLC graph of racemic 4an




HPLC graph of enantioenriched 4an




HPLC graph of racemic 4ba

475.46173 50.5246

1.8482 5.27246e4




HPLC graph of enantioenriched 4ba




[min] [min] [mAU*s] [mAU] %
----|-----|-----|------|------|
1 31.937 MF 2.7021 1.39181e5 858.45691 50.2613
2 50.030 FM 4.0753 1.37733e5 563.28522 49.7387

HPLC graph of racemic 4ca




Peak RetTime Type Width Height Area Area [mAU*s] # [min] [min] [mAU] 1 33.879 MM 2.4002 1085.09412 7.53462 1.0873 2 50.596 MM 3.9629 9.87165e4 415.16568

HPLC graph of enantioenriched 4ca




2 30.099 BB 0.9306 3.69286e4 601.21588 50.0692


HPLC graph of racemic 4da




HPLC graph of enantioenriched 4da




HPLC graph of racemic 4ea




HPLC graph of enantioenriched 4ea



HPLC graph of racemic 4ci




HPLC graph of enantioenriched 4ci



Peak	RetTime	Type	Width	Area	Height	Area	
#	[min]		[min]	[mAU*s]	[mAU]	용	
1	12.326	MM	1.2153	1.36443e4	187.11345	50.4274	
2	33.615	MM	1.9803	1.34130e4	112.88481	49.5726	

HPLC graph of racemic 5



HPLC graph of enantioenriched 5