

# **Multistep Mass-Remainder Analysis and its Application for Copolymer Blends**

Tibor Nagy<sup>1</sup>, Ákos Kuki<sup>1</sup>, Mahir Hashimov<sup>1,2</sup>, Miklós Zsuga<sup>1</sup>, Sándor Kéki<sup>1\*</sup>


<sup>1</sup>Department of Applied Chemistry, Faculty of Science and Technology, University of Debrecen, H-4032 Debrecen, Egyetem tér 1., Hungary

<sup>2</sup>University of Debrecen, Doctoral School of Chemistry, H-4032 Debrecen, Egyetem tér 1., Hungary.


\* Corresponding author: [keki.sandor@science.unideb.hu](mailto:keki.sandor@science.unideb.hu), fax: +36 52 518662; H-4032 Debrecen, Egyetem tér 1., HUNGARY

## *Table of content*

<b>FIGURE S1. MASS SPECTRUM OF RF551 COPOLYMER (SAMPLE 7.).....</b>	<b>2</b>
<b>FIGURE S2. MASS SPECTRUM OF PE3100 COPOLYMER (SAMPLE 8.).....</b>	<b>2</b>
<b>FIGURE S3. MASS SPECTRUM OF RPE1740 COPOLYMER (SAMPLE 9.).....</b>	<b>3</b>
<b>FIGURE S4. KMD PLOT OF EO/PO COPOLYMER BLEND (SAMPLE 6.) .....</b>	<b>3</b>



*Figure S1. Mass spectrum of RF551 copolymer (Sample 7.)*



*Figure S2. Mass spectrum of PE3100 copolymer (Sample 8.)*




Figure S3. Mass spectrum of RPE1740 copolymer (Sample 9.)




Figure S4. KMD plot of EO/PO copolymer blend (Sample 6.)