Supplementary Information

Superhydrophobic/superoleophilic surfaces by electroless nanoparticle deposition and perfluorinated polymer modification

Eugene B. Caldona[a], Hunter O. Brown[a,b], Dennis W. Smith, Jr.[a], and David O. Wipf[a*]

[a]Department of Chemistry and Marvin B. Dow Advanced Composites Institute, Mississippi State University, Mississippi State, MS 39762, United States
[b]Dave C. Swalm School of Chemical Engineering, Mississippi State University, Mississippi State, MS 39762, United States

*Corresponding author (DWipf@chemistry.msstate.edu)

Contents

Fig. S1. SEM images (inset: corresponding high magnification images) and EDX spectra for the uncoated and coated substrates after immersion for 7 d in 0.5 M NaCl solution. 2

Fig. S2. Screenshots illustrating the passage of (a) hexane, (b) petroleum ether, (c) gasoline, and (d) cooking oil through the **NFAg**-coated mesh. 2

Fig. S3. Sequential frames showing the separation of mixtures of water and (a) hexane, (b) petroleum ether, (c) gasoline, and (d) cooking oil. 3

Fig. S4. FTIR spectra of ACN mixed with (a) hexane, (b) petroleum ether, (c) gasoline, and (d) cooking oil containing different concentrations of water, with corresponding calibration curves of transmittance at 3600 cm^{-1} vs vol % H_2O. 3



Fig. S1. SEM images (inset: corresponding high magnification images) and EDX spectra for the uncoated and coated substrates after immersion for 7 d in 0.5 M NaCl solution.



Fig. S2. Screenshots illustrating the passage of (a) hexane, (b) petroleum ether, (c) gasoline, and (d) cooking oil through the **NFAg**-coated mesh.



Fig. S3. Sequential frames showing the separation of mixtures of water and (a) hexane, (b) petroleum ether, (c) gasoline, and (d) cooking oil.



Fig. S4. FTIR spectra of ACN mixed with (a) hexane, (b) petroleum ether, (c) gasoline, and (d) cooking oil containing different concentrations of water, with corresponding calibration curves of transmittance at 3600 cm^{-1} vs vol % H_2O.