

# Synthesis and chemical reactivity of tetrahydro[60]fullerene epoxides with both amino and aryl addends

Sisi Liang<sup>a</sup>, Liang Xu<sup>a</sup>, Liangbing Gan<sup>\*,ab</sup>

<sup>a</sup>Beijing National Laboratory for Molecular Sciences, Key Laboratory of Bioorganic

Chemistry and Molecular Engineering of the Ministry of Education, College of Chemistry

and Molecular Engineering, Peking University, Beijing 100871, China.

<sup>b</sup>State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic

Chemistry, Shanghai 200032, China.

## Table of Contents

Possible mechanism for the <b>10a</b> and <b>10b</b>	S2
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>3</b>	S3-S4
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>4</b>	S5-S6
<sup>1</sup> H NMR spectra for compound <b>5</b>	S7
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>6</b>	S8-S9
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>6a</b>	S10-S11
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>7</b>	S12-S13
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>8b</b>	S14-S15
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>8c</b>	S16-S17
<sup>1</sup> H and <sup>13</sup> C NMR and IR spectra for compound <b>9</b>	S18-S20
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>10a</b>	S21-S22
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>10b</b>	S23-S24
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>11</b>	S25-S26
<sup>1</sup> H NMR spectra for compound <b>13</b>	S27
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>14</b>	S28-S29
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>15</b>	S30-S31
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>16</b>	S32-S34
<sup>1</sup> H and <sup>13</sup> C NMR spectra for compound <b>17</b>	S35-S36

Possible mechanism for formation of **10a** and **10b**

Reference Huang, S. H.; Wang, F. D.; Gan, L. B.; Yuan, G.; Zhou, J.; Zhang, S. W. *Org. Lett.* **2006**, *8*, 277. In this work compound **M** was the only isolated product.













































































