

Supporting Information

Synthesis and characterization of $\text{NaNiF}_3 \cdot 3\text{H}_2\text{O}$: an unusual ordered variant of the ReO_3 type

Elena C. Gonzalo^{1,‡}, María Luisa Sanjuán², Markus Hoelzel³, M. Teresa Azcondo¹, Ulises Amador¹, Isabel Sobrados⁴, Jesús Sanz⁴, Flaviano García-Alvarado¹ and Alois Kuhn^{1,*}

¹ Universidad CEU San Pablo. Facultad de Farmacia, Departamento de Química y Bioquímica, Urbanización Montepríncipe, 28668 Boadilla del Monte, Madrid, Spain.

‡ Present address: CIC Energigune, Parque Tecnológico de Álava, 01510 Miñano, Spain

² Instituto de Ciencia de Materiales de Aragón (Universidad de Zaragoza-CSIC), Facultad de Ciencias, 50009 Zaragoza, Spain

³ Forschungsneutronenquelle Heinz Maier-Leibnitz (FRM II), D-85747 Garching, Germany

⁴ Instituto de Ciencia de Materiales de Madrid, CSIC, Spain




Figure S1. Rietveld analysis of the XRD patterns of $\text{NaNiF}_3 \cdot 3\text{H}_2\text{O}$ modelled with a cubic ($Pn-3$) $2a \times 2a \times 2a$ supercell. Bragg positions, calculated and observed profiles are depicted. A small amount of the secondary NaF phase was considered (lower vertical bars) in refinements.