SUPPORTING INFORMATION

Redox- and pH-sensitive polymeric micelles based on poly(β-amino ester)-grafted-disulfide methylene oxide poly(ethylene glycol) for anticancer drug delivery

Quang Nam Bui, † Yi Li, † Moon-Sun Jang, $^{\#}$ Dai Phu Huynh, § Jung Hee Lee, $^{\#}$ and Doo Sung Lee † , *

[†]Theranostic Macromolecules Research Center, School of Chemical Engineering, Sungkyunkwan University, Suwon 440-746, Republic of Korea

*Department of Radiology, Samsung Medical Center, Sungkyunkwan University School of Medicine and Center for Molecular and Cellular Imaging, Samsung Biomedical Research Institute, Seoul, 135-710, Republic of Korea

§Faculty of Materials Technology, National Key Lab. of Polymer and Composite Materials, HoChiMinh University of Technology, Vietnam National University, HoChiMinh City, Vietnam

Corresponding Author

^{*} E-mail address: dslee@skku.edu (D.S. Lee), Tel.: +82 31 290 7267, Fax: +82 31 292 8790.




Figure S1. ¹H-NMR spectra of MPEG-COOH (A), NHS-activated MPEG (B), MPEG-Cystamine (C)




Figure S2. GPC trace of PAE-g-DSMPEG.