

SUPPORTING INFORMATION

Targeted decationized polyplexes for siRNA delivery

*Novo^a, Kaori M. Takeda^b, Tamara Petteta^a, George R. Dakwar^c, Joep B. van den Dikkenberg^a,
Katrien Remaut^c, Kevin Braeckmans^{c,d}, Cornelius F. van Nostrum^a, Enrico Mastrobattista^a, Wim
E. Hennink^{a*}*

^aDepartment of Pharmaceutics, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, 3584 CG Utrecht, The Netherlands

^bDepartment of Materials Engineering, Graduate School of Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan

^cLaboratory for General Biochemistry and Physical Pharmacy, Faculty of Pharmacy, Ghent University, Ghent Research Group on Nanomedicines, Harelbekestraat 72, 9000 Ghent, Belgium

^dCentre for Nano- and Biophotonics, Ghent University, Harelbekestraat 72, 9000 Ghent, Belgium

* Corresponding author. Tel.: +31 30 253 6964; fax: +31 30 251 7839. E-mail address: w.e.hennink@uu.nl (W.E. Hennink).




Figure S1. Time dependent siRNA released from decationized pHP-PEG-FA polyplexes when incubated in human plasma or 20 mM HEPES buffer pH 7.4, as measured by FCS. Results are expressed as mean \pm SD (n=3).




Figure S2. Luciferase expression (a) and relative metabolic activity determined by XTT (b) of Skov3-luc cells upon treatment decationized pHP-PEG-FA polymer at different concentrations. XTT results were normalized to buffer treated samples. Results are expressed as mean \pm SD (n=5).




Figure S3. Relative metabolic activity determined by XTT of Skov3-luc cells upon treatment decationized pHP-PEG-FA polyplexes at different concentrations. Results were normalized to buffer treated samples. Results are expressed as mean \pm SD (n=5).