



Catalytic hydroamination of unactivated olefins using a Co catalyst for complex molecule synthesis

Hiroki Shigehisa, Natsumi Koseki, Nao Shimizu, Mayu Fujisawa, Makoto Niitsu, and Kou Hiroya*


Faculty of Pharmacy, Musashino University, 1-1-20 Shinmachi Nishitokyo-Shi, Tokyo, 202-8585, Japan

SUPPORTING INFORMATION


Part 2 ^1H , ^{13}C NMR, X-ray





S2





4a




S3




S4





S5




4d









4f




S8




4f




S9




4h




S10







4j




S12




4j




S13




4k




S14




4k




S15




4I




S16




S17




4m





S18





4m






S19





S20










S23




S24





S25




6d









6e





S28





6e




S29




S30






S31




6g









6h






S34




6h









6i




S37




6j





S38





6j





S39




S40





S41









S43





S44




S45






S46






60







S49




6p





S50





6p





S51





S52





S53





S54





S55





S56





S57




S58




S59





S60




13a







S62




S63




17





S64





17




S65




S66




S67




5b






5b






5c





S70





5c




S71




S72




S73




5e





S74





5e




S75




S76




S77




5j




S78




5j




S79




5k




S80




5k







5l




S82




5l




S83




5l'




S84




5l'




S85




5m




S86




5m




S87




7a





S88





7a









S90




S91




S92




S93





7d







S95





S96






7e






S98









S100




S101




S102




S103




7i





S104




7i







S106




7j




S107




7k




S108




7k







71




S110




7l




S111




S112





7m
diastereomer a




CGE09124a_2





NIT01039a_5






S115







S116

CGE09124b_2




NIT01039b_5






S118





S119



CGE09125a_2




CGE10033a_6




S120





S121





S122




S123













7p




S127





S128





10






S129







S131








S133




S134





S135




14a







S137






14b







14b









S140





S141





S142



S143



S144



S145

X-ray crystallography data for 14a





Table 1. Crystal data and structure refinement for CGE09128.

Identification code	CGE09128	
Empirical formula	C ₁₅ H ₁₃ N O	
Formula weight	223.26	
Temperature	296(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2(1)/c	
Unit cell dimensions	a = 7.7421(18) Å	α = 90°.
	b = 21.210(5) Å	β = 98.222(2)°.
	c = 7.4082(17) Å	γ = 90°.
Volume	1204.0(5) Å ³	
Z	4	
Density (calculated)	1.232 Mg/m ³	
Absorption coefficient	0.077 mm ⁻¹	
F(000)	472	
Crystal size	0.22 x 0.20 x 0.16 mm ³	
Theta range for data collection	1.92 to 26.37°.	
Index ranges	-9 ≤ h ≤ 9, -26 ≤ k ≤ 26, -9 ≤ l ≤ 9	
Reflections collected	12252	
Independent reflections	2470 [R(int) = 0.0305]	
Completeness to theta = 26.37°	99.8 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9877 and 0.9832	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2470 / 0 / 155	

Goodness-of-fit on F^2	1.025
Final R indices [$I > 2\sigma(I)$]	$R1 = 0.0417$, $wR2 = 0.1016$
R indices (all data)	$R1 = 0.0650$, $wR2 = 0.1136$
Largest diff. peak and hole	0.130 and -0.150 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for CGE09128. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
C(1)	9537(3)	2987(1)	8229(3)	66(1)
C(2)	10197(3)	3343(1)	6918(3)	62(1)
C(3)	9696(2)	3962(1)	6617(2)	51(1)
C(4)	8523(2)	4210(1)	7675(2)	42(1)
C(5)	7734(2)	4836(1)	7624(2)	40(1)
C(6)	7058(2)	5859(1)	6638(2)	43(1)
C(7)	7427(2)	6246(1)	8147(2)	52(1)
C(8)	6639(2)	6827(1)	8177(3)	62(1)
C(9)	5467(2)	7025(1)	6723(3)	66(1)
C(10)	5903(2)	6068(1)	5167(2)	56(1)
C(11)	5099(2)	6644(1)	5222(3)	66(1)
C(12)	6744(2)	4278(1)	9970(2)	52(1)
C(13)	7412(2)	4399(1)	11946(2)	68(1)
C(14)	7894(2)	3865(1)	9010(2)	46(1)
C(15)	8391(2)	3242(1)	9295(2)	58(1)
N(1)	7914(2)	5273(1)	6512(2)	45(1)
O(1)	6697(1)	4870(1)	8956(2)	52(1)

Table 3. Bond lengths [Å] and angles [°] for CGE09128.

C(1)-C(15)	1.380(3)
C(1)-C(2)	1.384(3)
C(1)-H(1)	0.9300
C(2)-C(3)	1.377(2)
C(2)-H(2)	0.9300
C(3)-C(4)	1.386(2)
C(3)-H(3)	0.9300
C(4)-C(14)	1.374(2)
C(4)-C(5)	1.459(2)
C(5)-N(1)	1.2613(19)
C(5)-O(1)	1.3600(18)
C(6)-C(10)	1.380(2)
C(6)-C(7)	1.383(2)
C(6)-N(1)	1.419(2)
C(7)-C(8)	1.377(2)
C(7)-H(7)	0.9300
C(8)-C(9)	1.371(3)
C(8)-H(8)	0.9300
C(9)-C(11)	1.371(3)
C(9)-H(9)	0.9300
C(10)-C(11)	1.375(2)
C(10)-H(10)	0.9300
C(11)-H(11)	0.9300
C(12)-O(1)	1.4608(19)
C(12)-C(14)	1.499(2)
C(12)-C(13)	1.503(2)
C(12)-H(12)	0.9800
C(13)-H(13A)	0.9600
C(13)-H(13B)	0.9600
C(13)-H(13C)	0.9600
C(14)-C(15)	1.385(2)
C(15)-H(15)	0.9300
C(15)-C(1)-C(2)	121.28(17)
C(15)-C(1)-H(1)	119.4
C(2)-C(1)-H(1)	119.4

C(3)-C(2)-C(1)	120.67(17)
C(3)-C(2)-H(2)	119.7
C(1)-C(2)-H(2)	119.7
C(2)-C(3)-C(4)	117.59(16)
C(2)-C(3)-H(3)	121.2
C(4)-C(3)-H(3)	121.2
C(14)-C(4)-C(3)	122.17(15)
C(14)-C(4)-C(5)	108.22(13)
C(3)-C(4)-C(5)	129.60(14)
N(1)-C(5)-O(1)	124.57(14)
N(1)-C(5)-C(4)	126.87(14)
O(1)-C(5)-C(4)	108.53(12)
C(10)-C(6)-C(7)	118.94(15)
C(10)-C(6)-N(1)	119.01(15)
C(7)-C(6)-N(1)	121.92(14)
C(8)-C(7)-C(6)	120.23(16)
C(8)-C(7)-H(7)	119.9
C(6)-C(7)-H(7)	119.9
C(9)-C(8)-C(7)	120.39(17)
C(9)-C(8)-H(8)	119.8
C(7)-C(8)-H(8)	119.8
C(11)-C(9)-C(8)	119.63(17)
C(11)-C(9)-H(9)	120.2
C(8)-C(9)-H(9)	120.2
C(11)-C(10)-C(6)	120.42(17)
C(11)-C(10)-H(10)	119.8
C(6)-C(10)-H(10)	119.8
C(9)-C(11)-C(10)	120.37(17)
C(9)-C(11)-H(11)	119.8
C(10)-C(11)-H(11)	119.8
O(1)-C(12)-C(14)	103.31(12)
O(1)-C(12)-C(13)	109.52(14)
C(14)-C(12)-C(13)	114.63(14)
O(1)-C(12)-H(12)	109.7
C(14)-C(12)-H(12)	109.7
C(13)-C(12)-H(12)	109.7
C(12)-C(13)-H(13A)	109.5
C(12)-C(13)-H(13B)	109.5

H(13A)-C(13)-H(13B)	109.5
C(12)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5
C(4)-C(14)-C(15)	119.93(16)
C(4)-C(14)-C(12)	109.05(14)
C(15)-C(14)-C(12)	130.99(15)
C(1)-C(15)-C(14)	118.33(17)
C(1)-C(15)-H(15)	120.8
C(14)-C(15)-H(15)	120.8
C(5)-N(1)-C(6)	120.30(13)
C(5)-O(1)-C(12)	110.73(12)

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for CGE09128. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
C(1)	73(1)	43(1)	74(1)	-3(1)	-15(1)	4(1)
C(2)	66(1)	55(1)	62(1)	-10(1)	2(1)	11(1)
C(3)	55(1)	50(1)	49(1)	-3(1)	7(1)	4(1)
C(4)	40(1)	43(1)	43(1)	0(1)	1(1)	-2(1)
C(5)	32(1)	48(1)	40(1)	0(1)	5(1)	-1(1)
C(6)	37(1)	42(1)	50(1)	7(1)	10(1)	-2(1)
C(7)	44(1)	52(1)	59(1)	0(1)	1(1)	1(1)
C(8)	59(1)	50(1)	76(1)	-10(1)	3(1)	0(1)
C(9)	59(1)	46(1)	92(2)	7(1)	8(1)	8(1)
C(10)	57(1)	55(1)	53(1)	5(1)	0(1)	0(1)
C(11)	58(1)	61(1)	72(1)	15(1)	-6(1)	8(1)
C(12)	38(1)	62(1)	56(1)	14(1)	10(1)	-6(1)
C(13)	61(1)	91(2)	54(1)	7(1)	15(1)	2(1)
C(14)	38(1)	48(1)	48(1)	6(1)	-4(1)	-7(1)
C(15)	58(1)	51(1)	62(1)	11(1)	-8(1)	-11(1)
N(1)	44(1)	45(1)	48(1)	4(1)	12(1)	2(1)
O(1)	46(1)	57(1)	57(1)	11(1)	20(1)	7(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for CGE09128.

	x	y	z	U(eq)
H(1)	9872	2567	8394	79
H(2)	10986	3163	6234	74
H(3)	10130	4204	5736	62
H(7)	8210	6113	9145	63
H(8)	6903	7087	9190	75
H(9)	4925	7415	6754	79
H(10)	5668	5817	4131	67
H(11)	4300	6776	4236	79
H(12)	5568	4097	9855	62
H(13A)	8572	4568	12056	102
H(13B)	7430	4010	12615	102
H(13C)	6661	4695	12431	102
H(15)	7964	3001	10183	70

Table 6. Torsion angles [°] for CGE09128.

C(15)-C(1)-C(2)-C(3)	-1.3(3)
C(1)-C(2)-C(3)-C(4)	0.2(3)
C(2)-C(3)-C(4)-C(14)	1.4(2)
C(2)-C(3)-C(4)-C(5)	-178.03(15)
C(14)-C(4)-C(5)-N(1)	-175.76(15)
C(3)-C(4)-C(5)-N(1)	3.7(3)
C(14)-C(4)-C(5)-O(1)	2.13(16)
C(3)-C(4)-C(5)-O(1)	-178.38(15)
C(10)-C(6)-C(7)-C(8)	0.5(2)
N(1)-C(6)-C(7)-C(8)	176.40(14)
C(6)-C(7)-C(8)-C(9)	0.8(3)
C(7)-C(8)-C(9)-C(11)	-0.9(3)
C(7)-C(6)-C(10)-C(11)	-1.7(2)
N(1)-C(6)-C(10)-C(11)	-177.72(15)
C(8)-C(9)-C(11)-C(10)	-0.3(3)
C(6)-C(10)-C(11)-C(9)	1.6(3)
C(3)-C(4)-C(14)-C(15)	-2.0(2)
C(5)-C(4)-C(14)-C(15)	177.57(14)
C(3)-C(4)-C(14)-C(12)	176.70(14)
C(5)-C(4)-C(14)-C(12)	-3.76(16)
O(1)-C(12)-C(14)-C(4)	3.89(16)
C(13)-C(12)-C(14)-C(4)	-115.21(15)
O(1)-C(12)-C(14)-C(15)	-177.64(15)
C(13)-C(12)-C(14)-C(15)	63.3(2)
C(2)-C(1)-C(15)-C(14)	0.8(3)
C(4)-C(14)-C(15)-C(1)	0.8(2)
C(12)-C(14)-C(15)-C(1)	-177.49(16)
O(1)-C(5)-N(1)-C(6)	2.5(2)
C(4)-C(5)-N(1)-C(6)	-179.96(13)
C(10)-C(6)-N(1)-C(5)	-120.85(17)
C(7)-C(6)-N(1)-C(5)	63.3(2)
N(1)-C(5)-O(1)-C(12)	178.39(14)
C(4)-C(5)-O(1)-C(12)	0.44(16)
C(14)-C(12)-O(1)-C(5)	-2.58(16)
C(13)-C(12)-O(1)-C(5)	119.99(14)

Symmetry transformations used to generate equivalent atoms:

X-ray crystallography data for 16




Table 1. Crystal data and structure refinement for CGE09105.

Identification code	CGE09105	
Empirical formula	C ₁₃ H ₁₉ N O ₂ S	
Formula weight	253.35	
Temperature	296(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2(1)/c	
Unit cell dimensions	a = 6.5204(11) Å	α = 90°.
	b = 26.602(4) Å	β = 101.144(2)°.
	c = 7.9704(13) Å	γ = 90°.
Volume	1356.4(4) Å ³	
Z	4	
Density (calculated)	1.241 Mg/m ³	
Absorption coefficient	0.229 mm ⁻¹	
F(000)	544	
Crystal size	0.18 x 0.10 x 0.06 mm ³	
Theta range for data collection	1.53 to 26.36°.	
Index ranges	-8 ≤ h ≤ 8, -33 ≤ k ≤ 33, -9 ≤ l ≤ 9	
Reflections collected	14091	
Independent reflections	2760 [R(int) = 0.0324]	
Completeness to theta = 26.36°	99.9 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9864 and 0.9599	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2760 / 0 / 159	
Goodness-of-fit on F ²	1.004	

Final R indices [I>2sigma(I)]	R1 = 0.0535, wR2 = 0.1223
R indices (all data)	R1 = 0.0632, wR2 = 0.1285
Largest diff. peak and hole	0.271 and -0.432 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for CGE09105. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
C(1)	3523(5)	4248(1)	1105(4)	74(1)
C(2)	5702(4)	4123(1)	2131(3)	52(1)
C(3)	7625(3)	4063(1)	5257(3)	42(1)
C(4)	7542(3)	3605(1)	6347(3)	38(1)
C(5)	9161(3)	3392(1)	7514(3)	44(1)
C(6)	8832(4)	2967(1)	8419(3)	48(1)
C(7)	10577(5)	2727(1)	9671(4)	69(1)
C(8)	7139(5)	4567(1)	1955(4)	76(1)
C(9)	5580(3)	3392(1)	6144(3)	42(1)
C(10)	7954(4)	4538(1)	6333(4)	65(1)
C(11)	6822(4)	2757(1)	8156(3)	54(1)
C(12)	5189(4)	2968(1)	7036(3)	53(1)
C(13)	6544(6)	3646(1)	1465(4)	85(1)
N(1)	5618(3)	4079(1)	3985(2)	44(1)
O(1)	2659(4)	3436(1)	3375(3)	109(1)
O(2)	2647(3)	4063(1)	5586(3)	100(1)
S(1)	3833(1)	3744(1)	4669(1)	60(1)

Table 3. Bond lengths [Å] and angles [°] for CGE09105.

C(1)-C(2)	1.532(4)
C(1)-H(1A)	0.9600
C(1)-H(1B)	0.9600
C(1)-H(1C)	0.9600
C(2)-N(1)	1.494(3)
C(2)-C(13)	1.519(4)
C(2)-C(8)	1.529(4)
C(3)-N(1)	1.493(3)
C(3)-C(4)	1.504(3)
C(3)-C(10)	1.520(3)
C(3)-H(3)	0.9800
C(4)-C(9)	1.381(3)
C(4)-C(5)	1.385(3)
C(5)-C(6)	1.382(3)
C(5)-H(5)	0.9300
C(6)-C(11)	1.401(3)
C(6)-C(7)	1.503(3)
C(7)-H(7A)	0.9600
C(7)-H(7B)	0.9600
C(7)-H(7C)	0.9600
C(8)-H(8A)	0.9600
C(8)-H(8B)	0.9600
C(8)-H(8C)	0.9600
C(9)-C(12)	1.382(3)
C(9)-S(1)	1.743(2)
C(10)-H(10A)	0.9600
C(10)-H(10B)	0.9600
C(10)-H(10C)	0.9600
C(11)-C(12)	1.370(3)
C(11)-H(11)	0.9300
C(12)-H(12)	0.9300
C(13)-H(13A)	0.9600
C(13)-H(13B)	0.9600
C(13)-H(13C)	0.9600
N(1)-S(1)	1.6419(17)
O(1)-S(1)	1.419(2)

O(2)-S(1)	1.439(2)
C(2)-C(1)-H(1A)	109.5
C(2)-C(1)-H(1B)	109.5
H(1A)-C(1)-H(1B)	109.5
C(2)-C(1)-H(1C)	109.5
H(1A)-C(1)-H(1C)	109.5
H(1B)-C(1)-H(1C)	109.5
N(1)-C(2)-C(13)	111.4(2)
N(1)-C(2)-C(8)	106.9(2)
C(13)-C(2)-C(8)	110.1(2)
N(1)-C(2)-C(1)	109.7(2)
C(13)-C(2)-C(1)	110.8(2)
C(8)-C(2)-C(1)	107.8(2)
N(1)-C(3)-C(4)	106.91(16)
N(1)-C(3)-C(10)	111.61(18)
C(4)-C(3)-C(10)	111.44(19)
N(1)-C(3)-H(3)	108.9
C(4)-C(3)-H(3)	108.9
C(10)-C(3)-H(3)	108.9
C(9)-C(4)-C(5)	118.66(19)
C(9)-C(4)-C(3)	113.68(18)
C(5)-C(4)-C(3)	127.64(18)
C(6)-C(5)-C(4)	120.7(2)
C(6)-C(5)-H(5)	119.7
C(4)-C(5)-H(5)	119.7
C(5)-C(6)-C(11)	118.9(2)
C(5)-C(6)-C(7)	121.4(2)
C(11)-C(6)-C(7)	119.8(2)
C(6)-C(7)-H(7A)	109.5
C(6)-C(7)-H(7B)	109.5
H(7A)-C(7)-H(7B)	109.5
C(6)-C(7)-H(7C)	109.5
H(7A)-C(7)-H(7C)	109.5
H(7B)-C(7)-H(7C)	109.5
C(2)-C(8)-H(8A)	109.5
C(2)-C(8)-H(8B)	109.5
H(8A)-C(8)-H(8B)	109.5

C(2)-C(8)-H(8C)	109.5
H(8A)-C(8)-H(8C)	109.5
H(8B)-C(8)-H(8C)	109.5
C(4)-C(9)-C(12)	122.3(2)
C(4)-C(9)-S(1)	109.16(16)
C(12)-C(9)-S(1)	128.58(17)
C(3)-C(10)-H(10A)	109.5
C(3)-C(10)-H(10B)	109.5
H(10A)-C(10)-H(10B)	109.5
C(3)-C(10)-H(10C)	109.5
H(10A)-C(10)-H(10C)	109.5
H(10B)-C(10)-H(10C)	109.5
C(12)-C(11)-C(6)	121.4(2)
C(12)-C(11)-H(11)	119.3
C(6)-C(11)-H(11)	119.3
C(11)-C(12)-C(9)	118.1(2)
C(11)-C(12)-H(12)	121.0
C(9)-C(12)-H(12)	121.0
C(2)-C(13)-H(13A)	109.5
C(2)-C(13)-H(13B)	109.5
H(13A)-C(13)-H(13B)	109.5
C(2)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5
C(3)-N(1)-C(2)	118.61(17)
C(3)-N(1)-S(1)	110.10(13)
C(2)-N(1)-S(1)	122.34(15)
O(1)-S(1)-O(2)	116.22(16)
O(1)-S(1)-N(1)	112.94(12)
O(2)-S(1)-N(1)	109.38(12)
O(1)-S(1)-C(9)	111.79(13)
O(2)-S(1)-C(9)	108.67(12)
N(1)-S(1)-C(9)	95.93(9)

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for CGE09105. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
C(1)	78(2)	76(2)	60(2)	14(1)	-5(1)	4(2)
C(2)	61(1)	49(1)	49(1)	6(1)	15(1)	4(1)
C(3)	29(1)	51(1)	49(1)	6(1)	9(1)	-3(1)
C(4)	33(1)	42(1)	42(1)	-3(1)	13(1)	-3(1)
C(5)	35(1)	52(1)	47(1)	-1(1)	11(1)	2(1)
C(6)	57(1)	49(1)	39(1)	-1(1)	13(1)	10(1)
C(7)	73(2)	74(2)	59(2)	15(1)	11(1)	20(1)
C(8)	86(2)	75(2)	73(2)	24(2)	27(2)	-9(2)
C(9)	38(1)	48(1)	40(1)	-1(1)	9(1)	-6(1)
C(10)	66(2)	50(1)	76(2)	-1(1)	3(1)	-13(1)
C(11)	74(2)	44(1)	46(1)	2(1)	18(1)	-8(1)
C(12)	55(1)	55(1)	49(1)	-2(1)	14(1)	-21(1)
C(13)	103(2)	77(2)	80(2)	-16(2)	30(2)	17(2)
N(1)	32(1)	54(1)	47(1)	7(1)	9(1)	-3(1)
O(1)	90(2)	138(2)	78(2)	36(1)	-32(1)	-70(2)
O(2)	49(1)	129(2)	135(2)	47(2)	54(1)	33(1)
S(1)	30(1)	83(1)	64(1)	21(1)	4(1)	-11(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for CGE09105.

	x	y	z	U(eq)
H(1A)	2623	3962	1100	110
H(1B)	3623	4331	-49	110
H(1C)	2957	4529	1620	110
H(3)	8785	4027	4646	51
H(5)	10482	3538	7690	53
H(7A)	11886	2872	9545	104
H(7B)	10599	2372	9453	104
H(7C)	10362	2782	10814	104
H(8A)	6647	4861	2449	114
H(8B)	7137	4626	767	114
H(8C)	8534	4491	2539	114
H(10A)	7995	4824	5603	98
H(10B)	9250	4515	7139	98
H(10C)	6824	4577	6936	98
H(11)	6593	2469	8755	65
H(12)	3856	2829	6880	63
H(13A)	7913	3577	2116	127
H(13B)	6623	3690	283	127
H(13C)	5629	3371	1574	127

Table 6. Torsion angles [°] for CGE09105.

N(1)-C(3)-C(4)-C(9)	-12.5(2)
C(10)-C(3)-C(4)-C(9)	109.7(2)
N(1)-C(3)-C(4)-C(5)	169.20(19)
C(10)-C(3)-C(4)-C(5)	-68.6(3)
C(9)-C(4)-C(5)-C(6)	2.0(3)
C(3)-C(4)-C(5)-C(6)	-179.8(2)
C(4)-C(5)-C(6)-C(11)	-1.0(3)
C(4)-C(5)-C(6)-C(7)	179.0(2)
C(5)-C(4)-C(9)-C(12)	-1.5(3)
C(3)-C(4)-C(9)-C(12)	180.0(2)
C(5)-C(4)-C(9)-S(1)	177.82(16)
C(3)-C(4)-C(9)-S(1)	-0.6(2)
C(5)-C(6)-C(11)-C(12)	-0.5(3)
C(7)-C(6)-C(11)-C(12)	179.5(2)
C(6)-C(11)-C(12)-C(9)	0.9(3)
C(4)-C(9)-C(12)-C(11)	0.1(3)
S(1)-C(9)-C(12)-C(11)	-179.12(18)
C(4)-C(3)-N(1)-C(2)	-127.21(19)
C(10)-C(3)-N(1)-C(2)	110.7(2)
C(4)-C(3)-N(1)-S(1)	20.8(2)
C(10)-C(3)-N(1)-S(1)	-101.26(19)
C(13)-C(2)-N(1)-C(3)	69.2(3)
C(8)-C(2)-N(1)-C(3)	-51.1(3)
C(1)-C(2)-N(1)-C(3)	-167.8(2)
C(13)-C(2)-N(1)-S(1)	-74.8(3)
C(8)-C(2)-N(1)-S(1)	164.92(18)
C(1)-C(2)-N(1)-S(1)	48.3(3)
C(3)-N(1)-S(1)-O(1)	-135.97(18)
C(2)-N(1)-S(1)-O(1)	10.7(2)
C(3)-N(1)-S(1)-O(2)	92.92(17)
C(2)-N(1)-S(1)-O(2)	-120.45(19)
C(3)-N(1)-S(1)-C(9)	-19.27(16)
C(2)-N(1)-S(1)-C(9)	127.35(17)
C(4)-C(9)-S(1)-O(1)	129.48(18)
C(12)-C(9)-S(1)-O(1)	-51.2(3)
C(4)-C(9)-S(1)-O(2)	-100.93(18)

C(12)-C(9)-S(1)-O(2)	78.4(2)
C(4)-C(9)-S(1)-N(1)	11.86(17)
C(12)-C(9)-S(1)-N(1)	-168.8(2)

Symmetry transformations used to generate equivalent atoms:

X-ray crystallography data for 18




Table 1. Crystal data and structure refinement for Shigehisa_01.

Identification code	Shigehisa_01	
Empirical formula	C ₁₄ H ₂₁ N O ₂ S	
Formula weight	267.38	
Temperature	296(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	Pbca	
Unit cell dimensions	a = 17.888(3) Å	$\alpha = 90^\circ$.
	b = 9.4477(14) Å	$\beta = 90^\circ$.
	c = 18.022(3) Å	$\gamma = 90^\circ$.
Volume	3045.7(8) Å ³	
Z	8	
Density (calculated)	1.166 Mg/m ³	
Absorption coefficient	0.208 mm ⁻¹	
F(000)	1152	
Crystal size	0.25 x 0.18 x 0.10 mm ³	
Theta range for data collection	2.26 to 26.37°.	
Index ranges	-22 ≤ h ≤ 22, -11 ≤ k ≤ 11, -22 ≤ l ≤ 22	
Reflections collected	30548	
Independent reflections	3118 [R(int) = 0.0270]	
Completeness to theta = 26.37°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9795 and 0.9499	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	3118 / 0 / 168	
Goodness-of-fit on F ²	0.939	

Final R indices [I>2sigma(I)]	R1 = 0.0407, wR2 = 0.1119
R indices (all data)	R1 = 0.0491, wR2 = 0.1205
Largest diff. peak and hole	0.320 and -0.316 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for Shigehisa_01. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
C(1)	1328(1)	3898(2)	1904(1)	44(1)
C(2)	1958(1)	4398(2)	2276(1)	44(1)
C(3)	2746(1)	4002(2)	2052(1)	50(1)
C(4)	2801(1)	3434(2)	1274(1)	50(1)
C(5)	3558(1)	2815(3)	1087(1)	68(1)
C(6)	1836(1)	5287(2)	2879(1)	52(1)
C(7)	1128(1)	5671(2)	3115(1)	56(1)
C(8)	514(1)	5157(2)	2732(1)	60(1)
C(9)	610(1)	4277(2)	2131(1)	56(1)
C(10)	1028(2)	6623(3)	3777(1)	83(1)
C(11)	1180(1)	3182(3)	-295(1)	64(1)
C(12)	1188(2)	4530(4)	-756(2)	122(1)
C(13)	1847(2)	2284(4)	-522(2)	100(1)
C(14)	471(2)	2368(5)	-408(2)	127(2)
N(1)	1241(1)	3672(2)	486(1)	52(1)
O(1)	1015(1)	1444(1)	1278(1)	63(1)
O(2)	2259(1)	2265(1)	1213(1)	51(1)
S(1)	1403(1)	2754(1)	1141(1)	46(1)

Table 3. Bond lengths [Å] and angles [°] for Shigehisa_01.

C(1)-C(2)	1.394(2)
C(1)-C(9)	1.393(2)
C(1)-S(1)	1.7549(18)
C(2)-C(6)	1.391(3)
C(2)-C(3)	1.513(2)
C(3)-C(4)	1.504(3)
C(3)-H(3A)	0.9700
C(3)-H(3B)	0.9700
C(4)-O(2)	1.473(2)
C(4)-C(5)	1.513(2)
C(4)-H(4)	0.9800
C(5)-H(5A)	0.9600
C(5)-H(5B)	0.9600
C(5)-H(5C)	0.9600
C(6)-C(7)	1.385(3)
C(6)-H(6)	0.9300
C(7)-C(8)	1.384(3)
C(7)-C(10)	1.506(3)
C(8)-C(9)	1.376(3)
C(8)-H(8)	0.9300
C(9)-H(9)	0.9300
C(10)-H(10A)	0.9600
C(10)-H(10B)	0.9600
C(10)-H(10C)	0.9600
C(11)-N(1)	1.486(2)
C(11)-C(14)	1.497(3)
C(11)-C(13)	1.520(3)
C(11)-C(12)	1.520(4)
C(12)-H(12A)	0.9600
C(12)-H(12B)	0.9600
C(12)-H(12C)	0.9600
C(13)-H(13A)	0.9600
C(13)-H(13B)	0.9600
C(13)-H(13C)	0.9600
C(14)-H(14A)	0.9600
C(14)-H(14B)	0.9600

C(14)-H(14C)	0.9600
N(1)-S(1)	1.4919(17)
O(1)-S(1)	1.4403(14)
O(2)-S(1)	1.6056(13)
C(2)-C(1)-C(9)	121.20(17)
C(2)-C(1)-S(1)	121.54(13)
C(9)-C(1)-S(1)	117.25(14)
C(1)-C(2)-C(6)	116.94(16)
C(1)-C(2)-C(3)	122.81(16)
C(6)-C(2)-C(3)	120.25(16)
C(4)-C(3)-C(2)	113.47(15)
C(4)-C(3)-H(3A)	108.9
C(2)-C(3)-H(3A)	108.9
C(4)-C(3)-H(3B)	108.9
C(2)-C(3)-H(3B)	108.9
H(3A)-C(3)-H(3B)	107.7
O(2)-C(4)-C(5)	106.41(15)
O(2)-C(4)-C(3)	107.11(14)
C(5)-C(4)-C(3)	113.89(17)
O(2)-C(4)-H(4)	109.8
C(5)-C(4)-H(4)	109.8
C(3)-C(4)-H(4)	109.8
C(4)-C(5)-H(5A)	109.5
C(4)-C(5)-H(5B)	109.5
H(5A)-C(5)-H(5B)	109.5
C(4)-C(5)-H(5C)	109.5
H(5A)-C(5)-H(5C)	109.5
H(5B)-C(5)-H(5C)	109.5
C(7)-C(6)-C(2)	122.74(18)
C(7)-C(6)-H(6)	118.6
C(2)-C(6)-H(6)	118.6
C(6)-C(7)-C(8)	118.75(18)
C(6)-C(7)-C(10)	120.5(2)
C(8)-C(7)-C(10)	120.7(2)
C(9)-C(8)-C(7)	120.37(18)
C(9)-C(8)-H(8)	119.8
C(7)-C(8)-H(8)	119.8

C(8)-C(9)-C(1)	120.00(18)
C(8)-C(9)-H(9)	120.0
C(1)-C(9)-H(9)	120.0
C(7)-C(10)-H(10A)	109.5
C(7)-C(10)-H(10B)	109.5
H(10A)-C(10)-H(10B)	109.5
C(7)-C(10)-H(10C)	109.5
H(10A)-C(10)-H(10C)	109.5
H(10B)-C(10)-H(10C)	109.5
N(1)-C(11)-C(14)	110.56(19)
N(1)-C(11)-C(13)	111.74(19)
C(14)-C(11)-C(13)	110.0(2)
N(1)-C(11)-C(12)	104.8(2)
C(14)-C(11)-C(12)	111.3(3)
C(13)-C(11)-C(12)	108.3(2)
C(11)-C(12)-H(12A)	109.5
C(11)-C(12)-H(12B)	109.5
H(12A)-C(12)-H(12B)	109.5
C(11)-C(12)-H(12C)	109.5
H(12A)-C(12)-H(12C)	109.5
H(12B)-C(12)-H(12C)	109.5
C(11)-C(13)-H(13A)	109.5
C(11)-C(13)-H(13B)	109.5
H(13A)-C(13)-H(13B)	109.5
C(11)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5
C(11)-C(14)-H(14A)	109.5
C(11)-C(14)-H(14B)	109.5
H(14A)-C(14)-H(14B)	109.5
C(11)-C(14)-H(14C)	109.5
H(14A)-C(14)-H(14C)	109.5
H(14B)-C(14)-H(14C)	109.5
C(11)-N(1)-S(1)	125.62(14)
C(4)-O(2)-S(1)	114.75(11)
O(1)-S(1)-N(1)	122.80(9)
O(1)-S(1)-O(2)	101.44(8)
N(1)-S(1)-O(2)	114.54(8)

O(1)-S(1)-C(1)	111.00(9)
N(1)-S(1)-C(1)	104.32(9)
O(2)-S(1)-C(1)	100.78(7)

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for Shigehisa_01. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
C(1)	41(1)	44(1)	47(1)	3(1)	2(1)	2(1)
C(2)	41(1)	42(1)	48(1)	9(1)	-1(1)	2(1)
C(3)	37(1)	49(1)	64(1)	7(1)	-1(1)	1(1)
C(4)	39(1)	44(1)	68(1)	8(1)	8(1)	1(1)
C(5)	44(1)	66(1)	93(2)	0(1)	17(1)	7(1)
C(6)	51(1)	52(1)	52(1)	2(1)	-7(1)	0(1)
C(7)	60(1)	58(1)	51(1)	-4(1)	1(1)	8(1)
C(8)	45(1)	72(1)	63(1)	-8(1)	8(1)	9(1)
C(9)	39(1)	67(1)	61(1)	-8(1)	0(1)	2(1)
C(10)	83(2)	95(2)	72(2)	-30(1)	-1(1)	16(2)
C(11)	61(1)	80(1)	50(1)	-11(1)	1(1)	2(1)
C(12)	185(4)	123(3)	58(2)	8(2)	-6(2)	20(3)
C(13)	86(2)	134(3)	80(2)	-42(2)	14(1)	14(2)
C(14)	76(2)	222(4)	83(2)	-59(2)	1(2)	-46(2)
N(1)	53(1)	51(1)	51(1)	-6(1)	-1(1)	5(1)
O(1)	54(1)	46(1)	88(1)	-2(1)	7(1)	-10(1)
O(2)	43(1)	39(1)	71(1)	1(1)	6(1)	3(1)
S(1)	41(1)	40(1)	56(1)	-2(1)	4(1)	0(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for Shigehisa_01.

	x	y	z	U(eq)
H(3A)	3063	4831	2095	60
H(3B)	2934	3293	2394	60
H(4)	2675	4180	918	60
H(5A)	3684	2099	1444	101
H(5B)	3928	3550	1098	101
H(5C)	3542	2403	600	101
H(6)	2248	5638	3135	62
H(8)	35	5408	2881	72
H(9)	196	3935	1875	67
H(10A)	1277	6217	4198	125
H(10B)	505	6723	3884	125
H(10C)	1238	7536	3672	125
H(12A)	775	5120	-613	183
H(12B)	1145	4293	-1272	183
H(12C)	1648	5028	-672	183
H(13A)	2300	2799	-430	150
H(13B)	1812	2060	-1040	150
H(13C)	1850	1424	-238	150
H(14A)	491	1504	-128	190
H(14B)	412	2152	-925	190
H(14C)	54	2926	-242	190

Table 6. Torsion angles [°] for Shigehisa_01.

C(9)-C(1)-C(2)-C(6)	0.0(3)
S(1)-C(1)-C(2)-C(6)	-179.32(13)
C(9)-C(1)-C(2)-C(3)	179.46(17)
S(1)-C(1)-C(2)-C(3)	0.1(2)
C(1)-C(2)-C(3)-C(4)	18.6(2)
C(6)-C(2)-C(3)-C(4)	-161.96(16)
C(2)-C(3)-C(4)-O(2)	-53.24(19)
C(2)-C(3)-C(4)-C(5)	-170.60(16)
C(1)-C(2)-C(6)-C(7)	0.3(3)
C(3)-C(2)-C(6)-C(7)	-179.18(18)
C(2)-C(6)-C(7)-C(8)	-0.4(3)
C(2)-C(6)-C(7)-C(10)	179.3(2)
C(6)-C(7)-C(8)-C(9)	0.3(3)
C(10)-C(7)-C(8)-C(9)	-179.4(2)
C(7)-C(8)-C(9)-C(1)	0.0(3)
C(2)-C(1)-C(9)-C(8)	-0.2(3)
S(1)-C(1)-C(9)-C(8)	179.19(16)
C(14)-C(11)-N(1)-S(1)	72.3(3)
C(13)-C(11)-N(1)-S(1)	-50.6(3)
C(12)-C(11)-N(1)-S(1)	-167.6(2)
C(5)-C(4)-O(2)-S(1)	-162.09(13)
C(3)-C(4)-O(2)-S(1)	75.75(16)
C(11)-N(1)-S(1)-O(1)	-48.1(2)
C(11)-N(1)-S(1)-O(2)	75.56(18)
C(11)-N(1)-S(1)-C(1)	-175.25(16)
C(4)-O(2)-S(1)-O(1)	-166.32(13)
C(4)-O(2)-S(1)-N(1)	59.26(15)
C(4)-O(2)-S(1)-C(1)	-52.07(14)
C(2)-C(1)-S(1)-O(1)	121.10(15)
C(9)-C(1)-S(1)-O(1)	-58.25(17)
C(2)-C(1)-S(1)-N(1)	-104.72(16)
C(9)-C(1)-S(1)-N(1)	75.93(17)
C(2)-C(1)-S(1)-O(2)	14.29(16)
C(9)-C(1)-S(1)-O(2)	-165.07(15)

Symmetry transformations used to generate equivalent atoms: