

Facile preparation of paclitaxel loaded silk fibroin nanoparticles for enhanced antitumor efficacy by locoregional drug delivery

Puyuan Wu^{a,‡}, Qin Liu^{a,‡}, Rutian Li^a, Jing Wang^b, Xu Zhen^b, Guofeng Yue^c, Huiyu Wang^d, Fangbo Cui^a, Fenglei Wu^d, Mi Yang^a, Xiaoping Qian^a, Lixia Yu^a, Xiqun Jiang^b, Baorui Liu^{a,}*

^a The Comprehensive Cancer Centre of Drum Tower Hospital, Medical School of Nanjing University, Clinical Cancer Institute of Nanjing University, 321 Zhongshan Rd, Nanjing 210008, China

^b Laboratory of Mesoscopic Chemistry and Department of Polymer Science, Engineering College of Chemistry & Chemical Engineering, Nanjing University, Nanjing 210093, China

^c Nanjing Medical University, 140 Hanzhong Rd, Nanjing 210029, China

^d The Comprehensive Cancer Center of Drum Tower Hospital Affiliated to Nanjing Medical University, 321 Zhongshan Rd, Nanjing 210008, China

Corresponding Author: *E-mail: baorui.liu@nju.edu.cn




Figure S1. Morphological changes of BGC-823 after exposed to PTX and PTX-SF NPs at 6400ng/ml PTX for 48h. Cell remained normal morphology (A) in SF group (B). Cells turned round in PTX (C) and PTX-SF NPs (D) groups.




Figure S2. H&E staining of organs in each experimental group ($\times 200$).