

Supporting Information

Combinatorial high-throughput screening for highly active Pd-Ir-Ce based ternary catalysts in electrochemical oxygen reduction reaction

*Sung Hyeon Park,^a Chang Hyuck Choi,^a Jae Kang Koh,^b Chanhoo Pak,^c Seon-ah Jin,^c
and Seong Ihl Woo^{a,b,*}*

^aDepartment of Chemical and Biomolecular Engineering, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Republic of Korea.

^bGraduate School of EEWS (WCU), Korea Advanced Institute of Science and Technology, Daejeon 305-701, Republic of Korea.

^cSamsung Advanced Institute of Technology, Nongseo-dong, Giheung-gu, Yongin-si, Gyeonggi-do, Republic of Korea.

*Corresponding author: siwoo@kaist.ac.kr




Figure S1. XRD patterns of all the catalyst spots for Pd-Ir-Ce combinatorial array between the ranges of 17.7 to 77.7°. The red arrows indicate the XRD patterns of the carbon paper (support). The lines 1 through 11 are as described in Figure 1a.




Figure S2. XRD patterns all the catalyst spots for Pd-Ir-Ce combinatorial array, arranged for every line for each coordinate.

