Chemical design of ^{99m}Tc-labeled probes for targeting osteogenic tumor region

Supporting Information

Mashiho Yanagi[£], Tomoya Uehara[£]*, Yukie Uchida[£], Sachiko Kiyota[£], Mai Kinoshita[£], Yusuke Higaki[£], Hiromichi Akizawa[¶], Hirofumi Hanaoka[£], Yasushi Arano[£,]

Affiliations and addresses of the authors:

[£]Graduate School of Pharmaceutical Sciences, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba, 263-8675, Japan.

[¶]Health Science University of Hokkaido, 1757 Kanazawa, Tobetsu-cho, Ishikari-gun, Hokkaido, 061-0293, Japan.

For correspondence or reprint contact to: Tomoya Uehara Ph.D. Department of Molecular Imaging and Radiotherapy, Graduate School of Pharmaceutical Sciences, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba, 260-8675 Japan. Tel: +81-43-226-2898, Fax: +81-43-226-2897; E-mail: tuehara@chiba-u.jp

Elemental analyses data of *N,N'*-bis-tert-butoxycarbonyl-S,S'-bis-triphenylmethyl-ethylene dicysteine [EC(Boc)$_2$(Trt)$_2$] and ethylene dicysteine oxorhenium(V) ($^{185/188}$Re-EC).

N,N'-Bis-tert-butoxycarbonyl-S,S'-bis-triphenylmethyl-ethylene dicysteine [EC(Boc)$_2$(Trt)$_2$]: Anal. C. H. N. ($C_{56}H_{60}N_2O_8S_2 \cdot 2H_2O$); calcd. 67.99, 6.52, 2.83, Found, 67.99, 6.44, 2.67.

Ethylene dicysteine oxorhenium(V) ($^{185/188}$Re-EC): Anal. C. H. N. ($C_8H_{13}N_2O_5ReS_2$ 1/4H_2O) Calc. 20.39, 2.87, 5.95, Anal. 20.65, 2.78, 5.67.