

Supplementary information


Three-Dimensional Scaffolds to Evaluate Tumor Associated Fibroblast-Mediated Suppression of Breast Tumor Specific T Cells

Vy Phan-Lai^{a,b}, Stephen J. Florczyk^b, Forrest M. Kievit^b, Kui Wang^b, Ekram Gad^a, Nora L. Disis^a, and Miqin Zhang^{b,*}


^aDepartment of Medicine, University of Washington, Seattle, Washington 98109

^bDepartment of Materials Science & Engineering, University of Washington, Seattle, Washington 98195


Supplemental Figures



Supplemental Figure 1. RFP expression in stably transfected MMC-RFP line. (a) Brightfield phase contrast image and (b) fluorescence image of MMC-RFP cell line; scale bar, 10 μm .



Supplemental Figure 2. Morphology of MMC, fibroblast or T cell cultures grown on CA scaffolds or 2D culture plates, visualized by SEM. Images of: (a) MMC, (b) fibroblasts, and (c) T cells cultured on CA (top row) or 2D (bottom row); 2000 \times magnification; scale bar, 10 μ m. MMC: thick arrow, T cell: thin arrow, fibroblast: thin dashed arrow. Data is representative of two experiments.



Supplemental Figure 3. Morphology of fibroblast co-cultures grown on CA scaffolds or 2D culture plates, visualized by SEM. Images of: (a) MMC and fibroblasts, and (b) p98 T cells and fibroblasts cultured on CA (top row) and 2D (bottom row); 2000 \times magnification; scale bar, 10 μ m. MMC: thick arrow, T cell: thin arrow, fibroblast: thin dashed arrow. Data is representative of three experiments.