

Supporting Information for

Anti-proliferative and Hypoglycemic Cucurbitane-type Glycosides from the Fruits of *Momordica charantia*

Ping-Chun Hsiao,^{†,‡} Chia-Ching Liaw,[†] Syh-Yuan Hwang,[§] Hui-Ling Cheng,[†] Li-Jie Zhang,[†]

Chien-Chang Shen,[†] Feng-Lin Hsu,^{*,‡} and Yao-Haur Kuo ^{*,†,||}

[†] Division of Herbal Drugs and Natural Products, National Research Institute of Chinese Medicine,
Taipei 112, Taiwan

[‡] Graduate Institute of Pharmacognosy, Taipei Medical University, Taipei 110, Taiwan

[§] Endemic Species Research Institute, Council of Agriculture, Nantou 552, Taiwan

^{||} Graduate Institute of Integrated Medicine, China Medical University, Taichung 404, Taiwan

Corresponding Author phone: Feng-Lin Hsu, Professor, Tel: +886-2-27361661, E-mail:
hsu320@tmu.edu.tw; Yao-Haur Kuo, Professor; Tel: +886-2-2820-1999 ext. 7061; Fax:
+886-2-2823-6150; E-mail: kuoyh@nricm.edu.tw.

List of Supporting Information

Figure S1. The ^1H spectrum of kuguaoside A (**1**) (400 HMz in pyridine- d_5)

Figure S2. The ^{13}C spectrum of kuguaoside A (**1**) (100 HMz in pyridine- d_5)

Figure S3. The ^1H spectrum of kuguaoside B (**2**) (400 HMz in pyridine- d_5)

Figure S4. The ^{13}C spectrum of kuguaoside B (**2**) (100 HMz in pyridine- d_5)

Figure S5. The ^1H spectrum of kuguaoside C (**3**) (400 HMz in pyridine- d_5)

Figure S6. The ^{13}C spectrum of kuguaoside C (**3**) (100 HMz in pyridine- d_5)

Figure S7. The ^1H spectrum of kuguaoside D (**4**) (400 HMz in pyridine- d_5)

Figure S8. The ^{13}C spectrum of kuguaoside D (**4**) (100 HMz in pyridine- d_5)




Figure S1. The ^1H spectrum of kuguaoside A (1) (400 HMz in pyridine-d_5)




Figure S2. The ¹³C spectrum of kuguaoside A (1) (100 HMz in pyridine-*d*₅)




Figure S3. The ^1H spectrum of kuguaoside B (2) (400 HMz in pyridine-d_5)




Figure S4. The ¹³C spectrum of kuguaoside B (2) (100 HMz in pyridine-*d*₅)




Figure S5. The ^1H spectrum of kuguaoside C (**3**) (400 HMz in pyridine- d_5)




Figure S6. The ¹³C spectrum of kuguaoside C (3) (100 HMz in pyridine-*d*₅)




Figure S7. The ^1H spectrum of kuguaoside D (**4**) (400 HMz in pyridine- d_5)




Figure S8. The ¹³C spectrum of kuguaoside D (**4**) (100 HMz in pyridine-*d*₅)