

Eudesmane-type Sesquiterpenoid and Guaianolides from *Kandelia candel* in a Screening Program for Compounds to Overcome TRAIL-resistance

Tomohiro Minakawa[†], Kazufumi Toume[†], Midori A. Arai[†], Samir Kumar Sadhu[‡], Firoj Ahmed[§], and Masami Ishibashi^{†*}

[†]Graduate School of Pharmaceutical Sciences, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8675, Japan, [‡]Pharmacy Discipline, Life Science School, Khulna University, Khulna-9208, Bangladesh, [§]Department of Pharmacy, Jagannath University, Dhaka-1100, Bangladesh

Supporting information

¹H NMR spectra of compounds **1-4**

Figure S1	¹ H NMR spectrum of 1 in CDCl ₃	P. S2
Figure S2	¹ H NMR spectrum of 2 in CDCl ₃	P. S3
Figure S3	¹ H NMR spectrum of 3 in CDCl ₃	P. S4
Figure S4	¹ H NMR spectrum of 4 in CDCl ₃	P. S5

Figure S1 ^1H NMR spectrum of **1** in CDCl_3




Figure S2 ^1H NMR spectrum of **2** in CDCl_3




Figure S3 ^1H NMR spectrum of **3** in CDCl_3




Figure S4 ^1H NMR spectrum of **4** in CDCl_3

