

Supporting Information

for

New $^{99m}\text{Tc}(\text{CO})_3$ Mannosylated Dextran Bearing

S-Derivatized Cysteine Chelator for Sentinel Lymph Node Detection

I. Pirmettis^{1*}, Y. Arano², T. Tsotakos¹, K. Okada², A. Yamaguchi², T. Uehara², M. Morais³, J. D. G. Correia³, I. Santos³, M. Martins⁴, S. Pereira⁴, P. Kyprianidou¹, C. Triantis¹, M. Pelecanou⁵, M. Papadopoulos¹

¹Institute of Radioisotopes and Radiodiagnostic Products, NCSR “Demokritos” 15310 Ag. Paraskevi, Athens, Greece

²Graduate School of Pharmaceutical Sciences, Chiba University, Chiba, Japan

³Unidade de Ciências Químicas e Radiofarmacêuticas, ITN, Estrada Nacional 10, 2686-953 Sacavém, Portugal

⁴ CICECO, Universidade de Aveiro, Portugal

⁵Institute of Biology, NCSR “Demokritos” 15310 Ag. Paraskevi, Athens, Greece.

Contents

Figure S1. ^1H - ^1H correlation spectrum (COSY, range F1 5.60 - 1.61, range F2 5.59 - 1.65) of DCM20 (**3**) in D_2O at 25 °C.

Figure S2. ^1H - ^{13}C correlation spectrum (HSQC, range δ_{H} 5.92-1.52, range δ_{C} 107.48 - 20.31.) of DCM20 (**3**) in D_2O at 25 °C.

Figure S3. ^1H - ^{13}C correlation spectrum (HSQC, range δ_{H} 5.51-1.55, range δ_{C} 106.26 - 19.32) of $\text{Re}(\text{CO})_3\text{-DC15}$ (**6**) in D_2O at 25 °C.

Figure S1.




Figure S2.




Figure S3.

