

## **Supporting Information**

### **Ipomotaosides A - D, Resin Glycosides from the Aerial Parts of *Ipomoea batatas* and their Inhibitory Activity against COX-1 and COX-2**

Kazuko Yoshikawa,\* Chiho Yagi, Hiroshi Hama, Masami Tanaka, Shigenobu Arihara, and Toshihiro Hashimoto


*Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-Cho, Tokushima 770-8514, Japan*

## Contents of Supporting Information


	page
<b>S1.</b> $^1\text{H}$ NMR spectrum (600 MHz) of Ipomotaoside A ( <b>1</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	4
<b>S2.</b> $^{13}\text{C}$ NMR spectrum (150 MHz) of Ipomotaoside A ( <b>1</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	5
<b>S3.</b> COSY spectrum (600 MHz) of Ipomotaoside A ( <b>1</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	6
<b>S4.</b> HMQC spectrum (600 MHz) of Ipomotaoside A ( <b>1</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	7
<b>S5.</b> HMBC spectrum (600 MHz) of Ipomotaoside A ( <b>1</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	8
<b>S6.</b> ROESY spectrum (600 MHz) of Ipomotaoside A ( <b>1</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	9
<b>S7.</b> $^1\text{H}$ NMR spectrum (600 MHz) of Ipomotaoside B ( <b>2</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	10
<b>S8.</b> $^{13}\text{C}$ NMR spectrum (150 MHz) of Ipomotaoside B ( <b>2</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	11
<b>S9.</b> COSY spectrum (600 MHz) of Ipomotaoside B ( <b>2</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	12
<b>S10.</b> HMQC spectrum (600 MHz) of Ipomotaoside B ( <b>2</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	13
<b>S11.</b> HMBC spectrum (600 MHz) of Ipomotaoside B ( <b>2</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	14
<b>S12.</b> ROESY spectrum (600 MHz) of Ipomotaoside B ( <b>2</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	15
<b>S13.</b> $^1\text{H}$ NMR spectrum (600 MHz) of Ipomotaoside C ( <b>3</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	16
<b>S14.</b> $^{13}\text{C}$ NMR spectrum (150 MHz) of Ipomotaoside C ( <b>3</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	17
<b>S15.</b> COSY spectrum (600 MHz) of Ipomotaoside C ( <b>3</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	18
<b>S16.</b> HMQC spectrum (600 MHz) of Ipomotaoside C ( <b>3</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	19
<b>S17.</b> HMBC spectrum (600 MHz) of Ipomotaoside C ( <b>3</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	20
<b>S18.</b> ROESY spectrum (600 MHz) of Ipomotaoside C ( <b>3</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	21
<b>S19.</b> $^1\text{H}$ NMR spectrum (600 MHz) of Ipomotaoside D ( <b>4</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	22
<b>S20.</b> $^{13}\text{C}$ NMR spectrum (150 MHz) of Ipomotaoside D ( <b>4</b> ) in $\text{C}_5\text{D}_5\text{N}$ .	23

<b>S21.</b>	COSY spectrum (600 MHz) of Ipomotaoside D ( <b>4</b> ) in C <sub>5</sub> D <sub>5</sub> N.	24
<b>S22.</b>	HMQC spectrum (600 MHz) of Ipomotaoside D ( <b>4</b> ) in C <sub>5</sub> D <sub>5</sub> N.	25
<b>S23.</b>	HMBC spectrum (600 MHz) of Ipomotaoside D ( <b>4</b> ) in C <sub>5</sub> D <sub>5</sub> N.	26
<b>S24.</b>	ROESY spectrum (600 MHz) of Ipomotaoside D ( <b>4</b> ) in C <sub>5</sub> D <sub>5</sub> N.	27


**S1.**  $^1\text{H}$  NMR spectrum (600 MHz) of Ipomotaoside A (**1**) in  $\text{C}_5\text{D}_5\text{N}$ .




S2.  $^{13}\text{C}$  NMR spectrum (150 MHz) of Ipomotaoside A (**1**) in  $\text{C}_5\text{D}_5\text{N}$ .




S3. COSY spectrum (600 MHz) of Ipomotaoside A (**1**) in C<sub>5</sub>D<sub>5</sub>N.




S4. HMQC spectrum (600 MHz) of Ipomotaoside A (**1**) in C<sub>5</sub>D<sub>5</sub>N.




S5. HMBC spectrum (600 MHz) of Ipomotaoside A (**1**) in C<sub>5</sub>D<sub>5</sub>N.




S6. ROESY spectrum (600 MHz) of Ipomotaoside A (**1**) in C<sub>5</sub>D<sub>5</sub>N.




S7.  $^1\text{H}$  NMR spectrum (600 MHz) of Ipomotaoside B (**2**) in  $\text{C}_5\text{D}_5\text{N}$ .




S8.  $^{13}\text{C}$  NMR spectrum (600 MHz) of Ipomotaoside B (**2**) in  $\text{C}_5\text{D}_5\text{N}$ .




S9. COSY spectrum (600 MHz) of Ipomotaoside B (**2**) in C<sub>5</sub>D<sub>5</sub>N.




S10. HMQC spectrum (600 MHz) of Ipomotaoside B (**2**) in C<sub>5</sub>D<sub>5</sub>N.




S11. HMBC spectrum (600 MHz) of Ipomotaoside B (**2**) in C<sub>5</sub>D<sub>5</sub>N.




S12. ROESY spectrum (600 MHz) of Ipomotaoside B (**2**) in C<sub>5</sub>D<sub>5</sub>N.




S13.  $^1\text{H}$  NMR spectrum (600Hz) of Ipomotaoside C (**3**) in  $\text{C}_5\text{D}_5\text{N}$ .




**S14.**  $^{13}\text{C}$  NMR spectrum (150 MHz) of Ipomotaoside C (**3**) in  $\text{C}_5\text{D}_5\text{N}$ .




**S15.** COSY spectrum (600 MHz) of Ipomotaoside C (**3**) in C<sub>5</sub>D<sub>5</sub>N.




S16. HMQC spectrum (600 MHz) of Ipomotaoside C (**3**) in C<sub>5</sub>D<sub>5</sub>N.




S17. HMBC spectrum (600 MHz) of Ipomotaoside C (**3**) in C<sub>5</sub>D<sub>5</sub>N.




S18. ROESY spectrum (600 MHz) of Ipomotaoside C (**3**) in C<sub>5</sub>D<sub>5</sub>N.




**S19.**  $^1\text{H}$  NMR spectrum (600 MHz) of Ipomotaoside D (**4**) in  $\text{C}_5\text{D}_5\text{N}$ .




S20.  $^{13}\text{C}$  NMR spectrum (150 MHz) of Ipomotaoside D (**4**) in  $\text{C}_5\text{D}_5\text{N}$ .




S21. COSY spectrum (600 MHz) of Ipomotaoside D (**4**) in C<sub>5</sub>D<sub>5</sub>N.



S22. HMQC spectrum (600 MHz) of Ipomotaoside D (**4**) in C<sub>5</sub>D<sub>5</sub>N.



S23. HMBC spectrum (600 MHz) of Ipomotaoside D (**4**) in C<sub>5</sub>D<sub>5</sub>N.



S24. ROESY spectrum (600 MHz) of Ipomotaoside D (**4**) in C<sub>5</sub>D<sub>5</sub>N.

