

Symbiopolyol: A VCAM-1 Inhibitor from a Symbiotic Dinoflagellate of the Jellyfish,

Mastigias papua

Novriyandi Hanif,[†] Osamu Ohno,^{†,‡} Makoto Kitamura,^{†,‡} Kaoru Yamada,^{†,‡} and Daisuke Uemura^{†,‡,*}

Department of Chemistry, Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8602, Japan, and Department of Biosciences and Informatics, Faculty of Science and Technology, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan

*To whom correspondence should be addressed. Tel/Fax: +81-45-566-1842. E-mail:

uemura@bio.keio.ac.jp

[†]Nagoya University

[‡]Present address: Faculty of Science and Technology, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan

List of Supporting Information

		Page
Figure S1	^1H NMR ($\text{CD}_3\text{OD}/\text{C}_5\text{D}_5\text{N}$ 2/1, 600 MHz) of symbiopolyol	3
Figure S2	^{13}C NMR ($\text{CD}_3\text{OD}/\text{C}_5\text{D}_5\text{N}$ 2/1, 150 MHz) of symbiopolyol	4
Figure S3	^1H - ^1H DQF-COSY NMR ($\text{CD}_3\text{OD}/\text{C}_5\text{D}_5\text{N}$ 2/1, 600 MHz) of symbiopolyol	5




Figure S1. ¹H NMR (CD₃OD/C₅D₅N 2/1, 600 MHz) of symbiopolyol (**1a**)




Figure S2. ¹³C NMR (CD₃OD/C₅D₅N 2/1, 150 MHz) of symbiopolyol (**1a**)




Figure S3. ¹H-¹H DQF-COSY NMR (CD₃OD/C₅D₅N 2/1, 600 MHz) of symbiopolylol (1a)