


## Highly Oxygenated Monoterpene Acylglucosides from *Spiraea cantoniensis*


Kaori Yoshida,<sup>†</sup> Atsuyuki Hishida,<sup>‡</sup> Osamu Iida,<sup>‡</sup> Keizo Hosokawa,<sup>§</sup> and Jun Kawabata<sup>\*,†</sup>

*Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan, Tsukuba Division, Research Center for Medicinal Plant Resources, National Institute of Biomedical Innovation, Tsukuba 305-0843, Japan, and Department of Nutritional Management, Faculty of Health Sciences, Hyogo University, Kakogawa 675-0195, Japan.*


<b>Contents</b>	<b>Page</b>
1) <b>Figure S1.</b> $^1\text{H}$ NMR spectrum of kodemarioside A ( <b>1</b> ; 500 MHz, acetone- $d_6$ )	2
2) <b>Figure S2.</b> $^{13}\text{C}$ NMR spectrum of kodemarioside A ( <b>1</b> ; 125 MHz, acetone- $d_6$ )	3
3) <b>Figure S3.</b> $^1\text{H}$ NMR spectrum of kodemarioside B ( <b>2</b> ; 500 MHz, acetone- $d_6$ )	4
4) <b>Figure S4.</b> $^{13}\text{C}$ NMR spectrum of kodemarioside B ( <b>2</b> ; 125 MHz, acetone- $d_6$ )	5
5) <b>Figure S5.</b> $^1\text{H}$ NMR spectrum of kodemarioside C ( <b>3</b> ; 500 MHz, acetone- $d_6$ )	6
6) <b>Figure S6.</b> $^{13}\text{C}$ NMR spectrum of kodemarioside C ( <b>3</b> ; 125 MHz, acetone- $d_6$ )	7
7) <b>Figure S7.</b> $^1\text{H}$ NMR spectrum of kodemarioside D ( <b>4</b> ; 500 MHz, acetone- $d_6$ )	8
8) <b>Figure S8.</b> $^{13}\text{C}$ NMR spectrum of kodemarioside D ( <b>4</b> ; 125 MHz, acetone- $d_6$ )	9
9) <b>Figure S9.</b> $^1\text{H}$ NMR spectrum of kodemarioside E ( <b>5</b> ; 500 MHz, acetone- $d_6$ )	10
10) <b>Figure S10.</b> $^1\text{H}$ NMR spectrum of kodemarioside F ( <b>6</b> ; 500 MHz, acetone- $d_6$ )	11
11) <b>Figure S11.</b> $^1\text{H}$ NMR spectrum of <b>7</b> (500 MHz, acetone- $d_6$ )	12
12) <b>Figure S12.</b> $^1\text{H}$ NMR spectrum of <b>8a</b> (500 MHz, chloroform- $d$ )	13
13) <b>Figure S13.</b> $^1\text{H}$ NMR spectrum of <b>8b</b> (500 MHz, chloroform- $d$ )	14
14) <b>Figure S14.</b> $^1\text{H}$ NMR spectrum of <b>9</b> (500 MHz, chloroform- $d$ )	15
15) <b>Figure S15.</b> $^1\text{H}$ NMR spectrum of <b>10R</b> (500 MHz, chloroform- $d$ )	16
16) <b>Figure S16.</b> $^1\text{H}$ NMR spectrum of <b>10S</b> (500 MHz, chloroform- $d$ )	17




**Figure S1.**  $^1\text{H}$  NMR spectrum of kodemarioside A (**1**; 500 MHz, acetone- $d_6$ )




**Figure S2.**  $^{13}\text{C}$  NMR spectrum of kodemarioside A (**1**; 125 MHz, acetone- $d_6$ )




**Figure S3.**  $^1\text{H}$  NMR spectrum of kodemarioside B (**2**; 500 MHz, acetone- $d_6$ )




**Figure S4.**  $^{13}\text{C}$  NMR spectrum of kodemarioside B (2; 125 MHz, acetone- $d_6$ )




**Figure S5.**  ${}^1\text{H}$  NMR spectrum of kodemarioside C (**3**; 500 MHz, acetone- $d_6$ )




**Figure S6.**  $^{13}\text{C}$  NMR spectrum of kodemarioside C (**3**; 125 MHz, acetone- $d_6$ )




**Figure S7.**  $^1\text{H}$  NMR spectrum of kodemarioside D (**4**; 500 MHz, acetone- $d_6$ )




**Figure S8.**  $^{13}\text{C}$  NMR spectrum of kodemarioside D (**4**; 125 MHz, acetone- $d_6$ )




**Figure S9.**  $^1\text{H}$  NMR spectrum of kodemarioside E (**5**; 500 MHz, acetone- $d_6$ )




**Figure S10.**  $^1\text{H}$  NMR spectrum of kodemarioside F (**6**; 500 MHz, acetone- $d_6$ )




**Figure S11.**  $^1\text{H}$  NMR spectrum of **7** (500 MHz, acetone- $d_6$ )




**Figure S12.**  $^1\text{H}$  NMR spectrum of **8a** (500 MHz, chloroform-*d*)




**Figure S13.**  $^1\text{H}$  NMR spectrum of **8b** (500 MHz, chloroform-*d*)



**Figure S14.**  ${}^1\text{H}$  NMR spectrum of **9** (500 MHz, chloroform-*d*)



**Figure S15.**  $^1\text{H}$  NMR spectrum of **10R** (500 MHz, chloroform-*d*)



**Figure S16.**  $^1\text{H}$  NMR spectrum of **10S** (500 MHz, chloroform-*d*)