

Combining Very Large Quadratic and Cubic Nonlinear Optical Responses in Extended, Tris-Chelate Metallochromophores with Six π -Conjugated Pyridinium Substituents

Benjamin J. Coe, John Fielden, Simon P. Foxon, Bruce S. Brunschwig, Inge Asselberghs, Koen Clays, Anna Samoc and Marek Samoc

Supporting Information

1. Additional Figures.....S2
2. Complete Reference.....S4

1. Additional Figures




Figure S1. MLCT and Stark spectra with calculated fits for complex salts **4** and **10** in external electric fields of 3.51×10^7 and 3.57×10^7 V m⁻¹, respectively. Top panel: absorption spectrum illustrating Gaussian curves used in data fitting; bottom panel: electroabsorption spectrum, experimental (blue) and fits (green) according to the Liptay equation.¹




Figure S2. MLCT and Stark spectra with calculated fits for complex salts **11** and **12B** in external electric fields of 3.57×10^7 and 5.26×10^7 V m⁻¹, respectively. Top panel: absorption spectrum illustrating Gaussian curves used in data fitting; bottom panel: electroabsorption spectrum, experimental (blue) and fits (green) according to the Liptay equation.¹

- (1) Liptay, W. In *Excited States*, Vol. 1; Lim, E. C., Ed.; Academic Press, New York, 1974, pp. 129–229).

2. Complete Reference

- (54) (d) Odom, S. A.; Webster, S.; Padilha, L. A.; Peceli, D.; Hu, H.-H.; Nootz, G.; Chung, S.-J.; Ohira, S.; Matichak, J. D.; Przhonska, O. V.; Kachkovski, A. D.; Barlow, S.; Brédas, J.-L.; Anderson, H. L.; Hagan, D. J.; Van Stryland, E. W.; Marder, S. R. *J. Am. Chem. Soc.* **2009**, *131*, 7510–7511.