

ELECTRONIC SUPPORTING INFORMATION

Preparation of Supramolecular Thiophene Host System Showing Solid-state Fluorescence by Using Chiral (1*R*,2*S*)-2-Amino-1,2-diphenylethanol

Naoki Shiota,[†] Takafumi Kinuta,[†] Tomohiro Sato,[‡] Nobuo Tajima,[§] Reiko Kuroda^{‡,⊥}, Yoshio Matsubara^{†,*}, and Yoshitane Imai^{†,*}

[†] Department of Applied Chemistry, Faculty of Science and Engineering, Kinki University, 3-4-1 Kowakae, Higashi-Osaka, Osaka 577-8502, Japan.

[‡] JST ERATO-SORST Kuroda Chiromorphology Team, 4-7-6, Komaba, Meguro-ku, Tokyo 153-0041, Japan.

[§] Graduate School of Pure and Applied Sciences, Tsukuba University, 1-1-1 Tenodai, Tsukuba, Ibaraki 305-8571, Japan

[⊥] Department of Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8902, Japan.

Correspondence to: y-imai@apch.kindai.ac.jp, y-matsu@apch.kindai.ac.jp

Figure ESI-1




Figure ESI-1. Solid-state fluorescence spectrum of complex **I**.

Figure ESI-2




Figure ESI-2. Solid-state fluorescence spectrum of complex **III**.