

Supporting Information



Bioactive Iridoid Glucosides from the Fruit of *Gardenia jasminoides*

Yang Yu,[†] Zuo-lei Xie, ^{||} Hao Gao,^{‡, †} Wei-wei Ma, ^{||} Yi Dai,^{‡, †} Ying Wang,^{‡, †}


Yi Zhong*, [§] and Xin-sheng Yao*, ^{†, ‡}

Department of Natural Products Chemistry, Shenyang Pharmaceutical University,
Shenyang 110016, P. R. China; Institute of Traditional Chinese Medicine & Natural
Products, College of Pharmacy, Jinan University, Guangzhou 510632, P. R. China;
Department of Biological Science and Biotechnology, Tsinghua University, Beijing
100084, P. R. China; Guangdong Province Key Laboratory of Pharmacodynamic
Constituents of TCM and New Drugs Research, Jinan University, Guangzhou
510632, P. R. China; Joekai Biotech. Co., Ltd., Beijing 100020, P.R. China


¹H- and ¹³C-NMR of compound 1 (CD₃OD)




¹H- and ¹³C-NMR of compound **2** (CD₃OD)




R=6-*O*-trans-p-coumaroyl- β -D-glucosyl




¹H- and ¹³C-NMR of compound **3** (CD₃OD)




R=6-*O*-trans-cinnamoyl- β -D-glucosyl




¹H- and ¹³C-NMR of compound 4 (DMSO-*d*₆)




R=*trans*-*p*-coumaroyl





¹H- and ¹³C-NMR of compound 4 (CD₃OD)





R=*trans*-p-coumaroyl




¹H- and ¹³C-NMR of compound 5 (CD₃OD)




¹H- and ¹³C-NMR of compound **6** (CD_3OD)




^1H - and ^{13}C -NMR of compound **7** (CD_3OD)




^1H - and ^{13}C -NMR of compound **8** (CD_3OD)





^1H - and ^{13}C -NMR of compound **9** (CD_3OD)



R=6-O-*trans*-sinapoyl- β -D-glucosyl

