

Flavonoid Glucuronides and a Chromone from the Aquatic Macrophyte *Stratiotes aloides*

Jürgen Conrad,^{*}† Bernhard Förster-Fromme,[†] Mihaela-Anca Constantin,[†] Vladimir Ondrus,[†]
Sabine Mika,[†] Fadime Mert-Balci,[†] Iris Klaiber,[‡] Jens Pfannstiel,[‡] Wolfgang Möller,[§] Harald Rösner,[§]
Karin Förster-Fromme,[⊥] Uwe Beifuss,[†]

[†]Institut für Chemie und [§]Zoologie, Universität Hohenheim, Garbenstrasse 30, D-70599 Stuttgart,
Germany

[‡]Institut für Physiologie, FG Biosensorik, Serviceeinheit des Life Science Centers, Universität
Hohenheim, August-v.Hartmann-Strasse 3, D-70599 Stuttgart, Germany

[⊥]Institut für Mikrobiologie, Universität Stuttgart, Allmandring 31, D-70569 Stuttgart, Germany

Figure S1. ¹H NMR spectrum and selective TOCSYs of **1** in pyridine-*d*₅

Figure S2. MS/MS-HCD-fragment ion spectrum of **1**.

Figure S1. ^1H NMR spectrum of **1** in pyridine- d_5 (A). Residual water resonance at δ 5.7 suppressed by presaturation. Selective 1D-TOCSYs on anomeric protons H-1'' (glucopyranosyl, B) and H-1''' (glucuronic acid moiety, C) displaying the individual ^1H spin systems of both carbohydrate moieties of **1**.




Figure S2. MS/MS-HCD-fragment ion spectrum of **1** (GlcA = glucopyranosiduronic acid; Glu = glucopyranosyl).

