

Supporting Information

References for the compounds tested for *in vitro* activity on growth of *X. fastidiosa*, which were not obtained from *Citrus sinensis* grafted on *C. limonia*:

The tested compounds rutaecarpin (35), N-methyl-4-methoxyquinolin-2-one (36), 5,7-dimethoxy-2,2-dimethyl-2*H*-1-benzopyran-6-propanoic acid (37), and guyanin (38) were isolated from *Hortia brasiliana*, *H. oreadica* (1); 3',4',5',5,7-pentamethoxyflavanone (39), 3',4',7,8-tetramethoxy-5,6-(2'',2''-dimethylpyrano)-flavone (43) from *Neoraputia magnifica* (2); 3-(1,1-dimethylallyl)-isoscopoletin (40), raunin (42) and gravelliferone methyl ether (41) from *Rauia resinosa* (Rutaceae) (3); odoratol (46), odoratone (47), gedunin (48) and catechin (44) were isolated from *Cedrela odorata* grafted on *Toona ciliata* (4); azadirachtin A (49) from *Azadirachta indica* (Meliaceae) (5), anacardic acid (45) from Anacardiaceae (5).

TABLE LEGENDS

Table 1. ^{13}C NMR chemical shifts for compounds **1-6, 8-10** and **18-20** (δ^a)

Footnotes:

Comparison of the ^{13}C NMR data of isolated compounds with those from literature: (6), (7), (8), (9). ^aSolvent, CDCl_3 , ^bPyridine- d_6 , ^cAcetone- d_6 , ^d assignments base on HSQC/HMBC. **6:** 7-*O*-Glc(6" \rightarrow 1''')rha: C-1 100.1, C-2 73.2, C-3 76.6, C-4 70.0, C-5 75.7, C-6 66.3; C-1 99.8, C-2 70.5, C-3 71.1, C-4 72.3, C-5 68.7, C-6 18.2.

Table 2. ^{13}C NMR chemical shifts for compounds **7, 11, 14, 25, 26, 31-33** (δ^a)

Footnotes:

Comparison of the ^{13}C NMR data of isolated compounds with those from literature: (10), (11), (12), (13), (14), (15), (16), (17); **31:** H-3': δ 3.85, d, $J = 6.9$; H-4': δ 4.98, d, $J = 6.9$; **32:** H-3': δ 3.69, d, $J = 8.6$; H-4': δ 4.70, d, $J = 8.6$. ^aSolvent, CDCl_3 , ^bAcetone- d_6 , ^c CD_3OD , ^d assignments base on HSQC/HMBC

Table3. ^{13}C NMR chemical shifts for compounds **12, 15, 17, 21, 23, 30, 34** (δ^a)

Footnotes:

Comparison of the ^{13}C NMR data of isolated compounds with those from literature: (16), (18), (19), (20), (21). ^aSolvent, CDCl_3 , ^bAcetone- d_6 , ^c assignments base on HSQC/HMBC

Literature Cited

- (1) Pasqualotto, V. G. Estudo fitoquímico e avaliação do potencial antimicrobial de espécies de *Hortia*: *H. oreadica*, *H. brasiliana* e *H. superba*. Master thesis, Universidade Federal de São Carlos, Programa de Pós-graduação em química, **2008**.
- (2) Tomazela, D. M.; Pupo, M. T.; Passador, E. A. P.; Silva, M. F. das G. F. da; Vieira, P. C.; Fernandes, J. B.; Rodrigues Fo, E.; Oliva, G.; Pirani, J. R. Pyrano chalcones and a flavone from *Neoraputia magnifica* and their *Trypanosoma cruzi* glycosomal glyceraldehyde-3-phosphate dehydrogenase-inhibitory act. *Phytochemistry*, **2000**, *55*, 643-651.
- (3) Velozo, E. S.; Oliveira, D. J.; Arruda, A. C.; Vieira, P. C.; Fernandes, J. B.; Silva, M. F. das G. F. da; Caracelli, I.; Zukerman-Schpector, J. Z. Rauianin, a new coumarin from *Rauia resinosa*. *Nat. Prod. Lett.*, **1997**, *9*, 237-244.
- (4) Silva, M.F. das G.F. da; Agostinho, S.M.M.; Paula, J.R. de; Oiano Neto, J.; Castro-Gamboa, I.; Rodrigues Fo., E.; Fernandes, J.B.; Vieira, P.C. Chemistry of *Toona ciliata* and *Cedrela odorata* graft (Meliaceae): Chemosystematic and Ecological Significance. *Pure Appl. Chem.*, **1999**, *71*, 1083-1087.
- (5) Leite, A. C.; Oliveira, C. G.; Godoy, M. P.; Bueno, F. C.; Oliveira, M. F. S. S.; Forim, M. R.; Fernandes, J. B.; Vieira, P. C.; Silva, M. F. das G. F. da; Bueno, O. C.; Pagnocca, F. C.; Hebling, M. J. A.; Bacci, M. J. Toxicity of *Cipadessa fruticosa* to the leaf-cutting ants *Atta sexdens rubropilosa* (Hymenoptera: Formicidae) and their symbiotic fungus. *Sociobiology*, **2005**, *46*, 17-26.

- (6) Chen, J.; Montanari, A. M.; Widmer, W. W. Two new polymethoxylated flavones, a class of compounds with potential anticancer activity, isolated from cold pressed Dancy tangerine peel oil solids. *J. Agric. Food Chem.*, **1997**, *45*, 364-368.
- (7) Machida, K.; Osawa, K. On the flavonoid constituents from the peels of *Citrus hassaku* Hort. Ex Tanaka. *Chem. Pharm. Bull.*, **1989**, *37*, 1092-1094.
- (8) Agrawal, P. K. Carbon-13 NMR of flavonoids. Elsevier Science, New York, 1989.
- (9) Chang, S. H. Flavonoids, coumarins and acridone alkaloids from the root bark of *Citrus limonia*. *Phytochemistry*, **1990**, *29*, 351-353.
- (10) Steck, W. New syntheses of demethylsuberosin, xanthyletin, (+/-)-decursidinol, (+)-marmesin, (-)-nodakenetin, (+/-)-decursin, and (+/-)-prantschimgin. *Can. J. Chem.*, **1971**, *49*, 2297-2300.
- (11) Patra, A.; Mitra, A. K. Carbon-13 NMR signals of some natural coumarins and their derivatives. *Org. Magn. Reson.*, **1981**, *17*, 222-224.
- (12) Ikeshiro, Y.; Mase, I.; Tomita, Y. Dihydropyranocoumarins from roots of *Peucedanum japonicum*. *Phytochemistry*, **1992**, *31*, 4303-4306.
- (13) Toro, M. J. U.; Muller, A. H.; Arruda, M. S. P.; Arruda, A. C. Alkaloids and coumarins from *Ticorea longiflora*. *Phytochemistry*, **1997**, *45*, 851-853.
- (14) Kong, L-Y; Yao, N-H; Niwa, M. Two new xanthyletin-type coumarins from *Peucedanum decursivum*. *Heterocycles*, **2000**, *53*, 2019-2025.
- (15) Ju-Ichi, M.; Inove, M.; Tsuda, R.; Shibukawa, N. Junosmarin, a new khellactone ester from *Citrus junos* Tanaka. *Heterocycles*, **1986**, *24*, 2777-2779.
- (16) Takemura, Y.; Kawaguchi, H.; Maki, S.; Ju-ichi, M.; Omura, M.; Ito, C.; Furukawa, H. Constituents of domestic *Citrus* plants. Part XXVI. Studies on the constituents

- of Yalaha. Structures of a new acridone alkaloid and two new coumarins. *Chem. Pharm. Bull.*, **1996**, *44*, 804-809.
- (17) Wu, T. S.; Furukawa, H. Acridone alkaloids. VII. Constituents of *Citrus sinensis* Osbeck var. *brasiliensis* Tanaka. Isolation and characterization of three new acridone alkaloids, and a new coumarin. *Chem. Pharm. Bull.*, **1983**, *31*, 901-906.
- (18) Burke, B. A.; Parkins, H. Coumarins from *Amyris balsamifera*. *Phytochemistry*, **1979**, *18*, 1073-1075.
- (19) Macias, F. A.; Massanet, G. M.; Rodriguez-Luiz, F.; Salva, J. ¹³C NMR of coumarins I: 3-(1,1-dimethylallyl) derivatives. *Magn. Reson. Chem.* **1989**, *27*, 705-707.
- (20) Zou, Y.; Lobera, M.; Snider, B. B. Synthesis of 2,3-dihydro-3-hydroxy-2-hydroxylalkylbenzofurans from epoxy aldehydes. One-step syntheses of brosimacutin G, vaginidiol, vaginol, smyrindiol, xanthoarnol, and avicenol A. Biomimetic syntheses of angelicin and psoralen. *J. Org. Chem.*, **2005**, *70*, 1761-1770.
- (21) Patra, A.; Misra, S. K.; Chaudhuri, S. K. Constituents of *Limonia acidissima*. Applications of two-dimensional NMR spectroscopy in structure elucidation. *J. Indian Chem. Soc.*, **1988**, *LXV*, 205-208.