

Ultrafast proton transfer to solvent: molecularity and intermediates from solvation- and diffusion-controlled regimes

J. L. Pérez-Lustres,^a F. Rodriguez-Prieto,^b M. Mosquera,*^b

T. A. Senyushkina,^a N. P. Ernsting,^a and S. A. Kovalenko*^a

^a*Institute for Chemistry, Humboldt University of Berlin*

Brook-Taylor-Strasse 2, 12489 Berlin (Germany)

E-mail: skovale@chemie.hu-berlin.de

^b*Physical Chemistry Department , Chemistry Faculty, University of Santiago*

Avda. Das Ciencias s/n, 15706 Santiago de Compostela (Spain)

E-mail: qfmmgfot@usc.es

Complete ref 26:

Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. P.; Head-Gordon, M.; Gonzalez, C.; Pople, J. A. *Gaussian 94*, revisions E.2 and B.2; Gaussian, Inc.: Pittsburgh, PA, 1995.