

Supporting Information

Low-Molecular-Weight Organo- and Hydrogelators

Based on Cyclo(*L*-Lys-*L*-Glu)

Huimin Geng^a, Lin Ye^a, Ai-ying Zhang^a, Jingbo Li^{a,b}, Zeng-guo Feng^{a,b}

^aSchool of Materials Science and Engineering, Beijing Institute of Technology, No. 5 South Street Zhongguancun, Beijing, 100081

^bBeijing Key Laboratory of Construction Tailorable Advanced Functional Materials and Green Applications, No. 5 South Street Zhongguancun, Beijing, 100081




Figure S1 Mass spectra of gelator **3** (a) and red xerogel aged for 10 days in acetone (b).




Figure S2 FT-IR spectra of fresh xerogel and red xerogel aged for 10 days.




Figure S3 XRD patterns of xerogel of **3** (a), **4** (b), **5** (c) and **6** (d) in n-butanol.




Figure S4 XRD patterns of fresh gel and red xerogel aged for 10 days in acetone of **3**.




Figure S5 ^1H NMR spectrum of **2** (400 MHz, $\text{DMSO}-d_6$).




Figure S6 ^{13}C NMR spectrum of **2** (100 MHz, $\text{DMSO}-d_6$).




Figure S7 ^1H NMR spectrum of **3** (400 MHz, $\text{DMSO}-d_6$).




Figure S8 ^{13}C NMR spectrum of **3** (100 MHz, $\text{DMSO}-d_6$).




Figure S9 ^1H NMR spectrum of **4** (400 MHz, $\text{DMSO}-d_6$).




Figure S10 ^{13}C NMR spectrum of **4** (100 MHz, $\text{DMSO}-d_6$).




Figure S11 ^1H NMR spectrum of **5** (400 MHz, $\text{DMSO}-d_6$).




Figure S12 ^{13}C NMR spectrum of **5** (100 MHz, $\text{DMSO}-d_6$).

