Supporting Information for:
Air-Initiated Radical Polymerization of Lithium Salts of
$ω$ -(Undecamethylcarba-closo-dodecaboran-1'-yl)alk-1-enes, $CH_2$ = $CH(CH_2)_{n-2}C(BMe)_{11}$ - $Li$
Kamesh Vyakaranam, Stefanie Körbe and Josef Michl*
Department of Chemistry and Biochemistry, University of Colorado, Boulder, 80309-0215
*To whom correspondence should be addressed; email: michl@eefus.colorado.edu
Department of Chemistry and Biochemistry, University of Colorado, Boulder, 80309-0215

#### **Supporting Information Table of Contents**

H NMR spectrum in acetone: poly-5[2] prepared with air initiation	S1
H NMR spectrum in acetone: poly- <b>6</b> [2] prepared with air initiation	S2
H NMR spectrum in acetone: poly-7[2] prepared with air initiation	S3
H NMR spectrum in acetone: poly-5[2] prepared with AIBN initiation	S4
H NMR spectrum in acetone: poly-6[2] prepared with AIBN initiation	S <b>5</b>
H NMR spectrum in acetone: poly-7[2] prepared with AIBN initiation	S6
H NMR spectrum in acetone: poly-5[2] prepared with DTBP initiation	S <b>7</b>
H NMR spectrum in acetone: poly-6[2] prepared with DTBP initiation	S8
H NMR spectrum in acetone: poly-7[2] prepared with DTBP initiation	S9
<sup>13</sup> C NMR spectrum in acetone: poly- <b>5</b> [2] prepared with air initiation	S10
<sup>13</sup> C NMR spectrum in acetone: poly- <b>6</b> [2] prepared with air initiation	S11
<sup>13</sup> C NMR spectrum in acetone: poly- <b>7</b> [2] prepared with air initiation	S12
<sup>13</sup> C NMR spectrum in acetone: poly- <b>5</b> [2] prepared with DTBP initiation	S13
<sup>13</sup> C NMR spectrum in acetone: poly- <b>6</b> [2] prepared with DTBP initiation	S14
<sup>13</sup> C NMR spectrum in acetone: poly-7[2] prepared with DTBP initiation	S15
<sup>13</sup> C NMR spectrum in acetone: poly- <b>5</b> [2] prepared with AIBN initiation	S16
<sup>13</sup> C NMR spectrum in acetone: poly- <b>6</b> [2] prepared with AIBN initiation	S17
<sup>13</sup> C NMR spectrum in acetone: poly- <b>6</b> [2] prepared with AIBN initiation	S18

#### <sup>1</sup>H NMR spectrum in acetone: poly-5[2] prepared with air initiation

# after trichloroacetyl isocyanate treatment original polymer NH-Me<sub>11</sub> Solvent 6 5 Chemical Shift (ppm)

2.0


1.5 1.0 Chemical Shift (ppm)

1.0


0.5

3.0


### <sup>1</sup>H NMR spectrum in acetone: poly-6[2] prepared with air initiation




#### <sup>1</sup>H NMR spectrum in acetone: poly-7[2] prepared with air initiation




#### <sup>1</sup>H NMR spectrum in acetone: poly-5[2] prepared with AIBN initiation




#### <sup>1</sup>H NMR spectrum in acetone: poly-6[2] prepared with AIBN initiation




## <sup>1</sup>H NMR spectrum in acetone: poly-7[2] prepared with AIBN initiation




#### <sup>1</sup>H NMR spectrum in acetone: poly-5[2] prepared with DTBP initiation




#### <sup>1</sup>H NMR spectrum in acetone: poly-6[2] prepared with DTBP initiation




#### <sup>1</sup>H NMR spectrum in acetone: poly-7[2] prepared with DTBP initiation




#### <sup>13</sup>C NMR spectrum in acetone: poly-5[2] prepared with air initiation




### <sup>13</sup>C NMR spectrum in acetone: poly-6[2] prepared with air initiation




# <sup>13</sup>C NMR spectrum in acetone: poly-7[2] prepared with air initiation




#### <sup>13</sup>C NMR spectrum in acetone: poly-5[2] prepared with DTBP initiation




#### <sup>13</sup>C NMR spectrum in acetone: poly-6[2] prepared with DTBP initiation




### <sup>13</sup>C NMR spectrum in acetone: poly-7[2] prepared with DTBP initiation



#### <sup>13</sup>C NMR spectrum in acetone: poly-5[2] prepared with AIBN initiation



#### <sup>13</sup>C NMR spectrum in acetone: poly-6[2] prepared with AIBN initiation



### <sup>13</sup>C NMR spectrum in acetone: poly-7[2] prepared with AIBN initiation

