

Anal. Chem., 1998, 70(18), 3906-3911, DOI:10.1021/ac971327u

Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at http://pubs.acs.org/page/copyright/permissions.html






Figure 2(B and C). Variation of enrichment factor, E_e , with extracted sample volume for the chloro-s-triazines (0.40 ppm each) at two concentrations of sulphuric acid as an acceptor solution, atrazine (B), terbuthylazine (C), (\bullet) 0.2 M H₂SO₄ (pH \sim 0.7 (\blacksquare) 1.0 M H₂SO₄ (pH \sim 0.0)




Figure 5(B and C). Variation of enrichment factor with extracted sample volume for chloro-s-triazines at different ionic strength of the donor solution, atrazine (B), terbuthylazine (C), (♦) 0.23, (■) 0.69, (♦) 1.59, (o) 3.1. The acceptor solution contained 1.0 M sulphuric acid.