JOURNAL OF THE AMERICAN CHEMICAL SOCIETY
J. Am. Chem. Soc., 1997, 119(17), 4050-4058, DOI:10.1021/ja963631v

Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at http://pubs.acs.org/page/copyright/permissions.html

ACS Publications
MOST TRUSTED. MOST CITED. MOST READ.

Copyright © 1997 American Chemical Society

Figure S1

^{1}H NMR spectra obtained at 15 min. intervals during exchange of the *syn*(O)- and *anti*(O)-N*H* protons in $[Co(Mecyclen)(O_2C_2O_2)]^+$, D_2O solution, pD = 2.81, 25.0°C. The non-decaying signal (7.0 ppm) represents the *ap*-N*H* proton.

