

- 1 Eid, C. N.; Konopelski, J. P. *Tetrahedron Lett.* **1991**, 32, 461.
- 2 Meng, D.; Sorensen, E. J.; Bertinato, P.; Danishefsky, S. J. *J. Org. Chem.* **1996**, 61, 7998.
- 3 Keck, G. E.; Tarbet, K. H.; Geraci, L. S. *J. Am. Chem. Soc.* **1993**, 115, 8467. The allylic alcohol could also be prepared according to the general procedure outlined in the following references: (a) Racherla, U. S.; Brown, H. C. *J. Org. Chem.* **1991**, 56, 401. (b) Yang, Z.; He, Y.; Vourloumis, D.; Vallberg, H.; Nicolaou, K. C. *Angew. Chem., Int. Ed. Engl.*, **1997**, 36, 166.
- 4 Prepared according to the procedure outlined in an earlier Danishefsky synthesis of the epothilones: Meng, D.; Sorensen, E. J.; Bertinato, P.; Danishefsky, S. J. *J. Org. Chem.* **1996**, 61, 7998.
- 5 The oxidation procedure described here may also be accomplished with NaIO₄ as outlined in a previous Danishefsky synthesis of the epothilones: Meng, D.; Bertinato, P.; Balog, A.; Su, D.-S.; Kamenecka, T.; Sorensen, E. J.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1997**, 119, 10073
- 6 (a) Stork, G.; Zhao, K. *Tetrahedron Lett.* **1989**, 30, 2173. (b) Stork, G.; Zhao, K. *J. Am. Chem. Soc.* **1990**, 112, 5875. (c) Chen, J.; Wang, T.; Zhao, K. *Tetrahedron Lett.* **1994**, 35, 2827.
- 7 The substrate was prepared by treating the Overman alcohol (see reference 8) with TESOTf and 2,6-lutidine in CH₂Cl₂ at -78 °C for 30 min.
- 8 Aldehydes **49a** and **49b** were readily prepared according to the procedure outlined in: Lin, N.-H.; Overman, L. E.; Rabinowitz, M. H.; Robinson, L. A.; Sharp, M. J.; Zablocki, J. *J. Am. Chem. Soc.* **1996**, 118, 9062.
- 9 Roush, W. R.; Hoong, L. K.; Palmer, M. A. J.; Straub, J. A.; Palkowitz, A. D. *J. Org. Chem.* **1990**, 55, 4117.