

Supporting Information

Highly Efficient $\text{B}(\text{C}_6\text{F}_5)_3$ -Catalyzed Hydrosilylation of Olefins

Michael Rubin, Todd Schwier, and Vladimir Gevorgyan*

Department of Chemistry, University of Illinois at Chicago
845 West Taylor Street, Chicago, Illinois 60607-7061


^1H and ^{13}C charts of new compound **3n** (Total number of pages: 3).

Current Data Parameters
 NAME mr0596
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20010813
 Time 16.46
 INSTRUM spect
 PROBHD 5 mm BBO BB-1
 PULPROG zg
 TD 16384
 SOLVENT CDCl3
 NS 8
 DS 2
 SMH 4789.272 Hz
 FIDRES 0.292314 Hz
 AQ 1.7105396 sec
 RG 32
 DM 104.400 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec
 P1 5.50 usec
 SF01 400.1342000 MHz
 NUC1 ¹H
 PL1 -3.00 dB

F2 - Processing parameters
 SI 8192
 SF 400.1320090 MHz
 WDW EM
 SSB 0
 LB 0.10 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 F1P 10.100 ppm
 F1 4041.33 Hz
 F2P -0.100 ppm
 F2 -40.01 Hz
 PPMCM 0.51000 ppm/cm
 HZCM 204.06735 Hz/cm



Carbon-13 standard parameters, BBO probe

