




5NIM/N098-3-19-D-GLUCURONIC-HCL

D:\B8425NIM.GXD;1
COMNT B8425NIM/N098-3-19-D-GLUCURONI
DATIM 20-MAR-98 13:21:01
OENUC 1H
EXMOD HMG
OBFRO 399.65 MHz
OBSET 124.00 KHz
OBFIN 10500.0 Hz
POINT 16384
FREQU 8000.0 Hz
SCANS 16
ACQTM 2.048 sec
PD 4.952 sec
PWL 5.5 us
IRNUC 1H
CTEMP 24.0 c
SLVNT D2O
EXREF 4.65 ppm
BF 0.12 Hz
RGAIN 24




9NIM/L-IDURONIC (LARGE)

2059.18
 2052.84
 1962.99
 1961.04
 1871.20
 1862.41
 1858.99
 1659.28
 1655.38
 1652.93
 1648.54
 1633.40
 1627.06
 1620.71
 1465.92
 1408.79
 1401.47
 1395.12
 1387.80
 1330.67
 1318.46
 1314.56
 1304.79
 1300.88
 1290.14
 1284.28
 1279.89
 1276.47
 1272.56
 1268.66




D:\B8397NIM.GXD;1
 B8379NIM/L-IDURONIC (LARGE)
 17-MAR-98 16:47:51
 1H
 EXMOD HMG
 399.65 MHz
 124.00 KHz
 10500.0 Hz
 16384
 8000.0 Hz
 16
 2.048 sec
 4.952 sec
 5.5 us
 23.8 c
 4.65 ppm
 0.12 Hz
 27





NIM/NO98-4-21-L-D-MANN-TFA-HCL-SALT

DFILE D:\B8669NIM.GXD;1
 COMNT B8669NIM/NO98-4-21-L-D-MANN-TFA
 DATIM 22-APR-98 17:34:24
 EXMOD 1H
 OBNUC 1H
 EXMOD HMG
 OBFREQ 399.65 MHz
 OBSET 124.00 KHz
 OBFIN 10500.0 Hz
 POINT 16384
 FREQU 8000.0 Hz
 SCANS 128
 ACQTM 2.048 sec
 PD 4.952 sec
 PW1 5.5 us
 IRNUC 1H
 CTMP 23.3 C
 SLVNT D2O
 EXREF 4.65 ppm
 BF 0.12 Hz
 RGAIN 31



9NIM/NO98-4-8-H-(GULURONIC(?))-TFA-HCL

DFILE D:\B8579NIM.GXD;1
 COMNT B8579NIM/NO98-4-8-H-(GULURONIC
 DATIM 09-APR-98 15:19:03
 OBNUC 1H
 EXMOD HMG
 OBFREQ 399.65 MHz
 OBSET 124.00 kHz
 OBFIN 10500.0 Hz
 POINT 16384
 FREQU 8000.0 Hz
 SCANS 32
 ACQTM 2.048 sec
 PD 4.952 sec
 FW 5.5 us
 IRNUC 1H
 CTMP 23.9 c
 SLVNT D2O
 EXREF 4.65 ppm
 BF 0.12 Hz
 RGAIN 29





6NIM/NPHT-BOC-DI-MOM-RFH



DFILE
 COMNT
 DATIM
 OBNUC
 EXMOD
 OBFRQ
 OBSET
 OBFIN
 POINT
 FREQU
 SCANS
 ACQTM
 PD
 PW1
 IRNUC
 CTMP
 SLVNT
 EXREF
 BF
 RGAIN

D:\B6786NIM.GXD;1
 B6786NIM/NPHT-BOC-DI-MOM-RF
 14-AUG-97 16:48:02
 1H
 NON

399.65 MHz
 124.00 KHz
 10500.0 Hz
 16384
 8000.0 Hz
 16
 2.048 sec
 4.952 sec
 5.4 us
 1H
 CDCL3
 25.0 C
 0.00 ppm
 0.12 Hz
 16




D:\B6506NIM.GXD;1
 B6506NIM/NO97-6-26-L
 26-JUN-97 16:12:10
 1H
 NON
 399.65 MHz
 124.00 KHz
 10500.0 Hz
 16384
 8000.0 Hz
 16
 2.048 sec
 4.952 sec
 5.4 us
 1H
 24.5 c
 CDCL3
 0.00 ppm
 0.12 Hz
 13
 RGAIN




4NIM/D-MANN-MOM-PHT-BOC-CH2OH-L

DFILE
COMPT
DATIM
OBNUC
EXMOD
OBFRO
OBSET
OBFIN
POINT
FREOU
SCANS
ACQTM
PD
FWI
IRNUC
CTEMP
SLVNT
EXREF
BF
RGAIN


D:\B7614NIM.GXD;1
B7614NIM/D-MANN-MOM-PHT-BOC-CH
28-NOV-97 20:19:36
1H
NON
399.65 MHz
124.00 KHz
10500.0 Hz
16384
8000.0 Hz
16
2.048 sec
4.952 sec
5.4 us
1H
CDCL3
24.6 c
0.00 ppm
0.12 Hz
15



2NIM/L-GULO-MOM-PHT-BOC-CH2OH-H



D:\B7612NIM.GXD;1
 COMNT B7612NIM/L-GULO-MOM-PHT-BOC-C
 DATIM 28-NOV-97 19:38:54
 OBNUC 1H
 EXMOD NON
 OBFRQ 399.65 MHz
 OBSET 124.00 KHz
 OBFIN 10500.0 Hz
 POINT 16384
 FREQU 8000.0 Hz
 SCANS 16
 ACQTM 2.048 sec
 PD 4.952 sec
 PW1 5.4 us
 TRNUC 1H
 CTEMP 24.8 c
 SLVNT CDCL3
 EXREF 0.00 ppm
 BF 0.12 Hz
 RGAIN 16




Flexible Synthesis and Biological Activity of Uronic Acid-Type *Gem*-diamine 1-*N*-Iminosugars: A New Family of Glycosidase Inhibitors

Yoshio Nishimura,* Eiki Shitara, Hayamitsu Adachi, and Tomio Takeuchi

Institute of Microbial Chemistry, 3-14-23 Kamiosaki, Shinagawa-ku, Tokyo 141-0021, Japan

X-ray Data of Compound 24

nm 5.10/10



*Experimental*Data Collection

A colorless prism crystal of $C_{23}H_{30}N_2O_8$ having approximate dimensions of 0.35 x 0.20 x 0.20 mm was mounted on a glass fiber. All measurements were made on a Rigaku AFC7R diffractometer with graphite monochromated Cu-K α radiation and a rotating anode generator.

Cell constants and an orientation matrix for data collection, obtained from a least-squares refinement using the setting angles of 25 carefully centered reflections in the range $57.31 < 2\theta < 59.93^\circ$ corresponded to a primitive orthorhombic cell with dimensions:

$$\begin{aligned}a &= 11.610(2) \text{ \AA} \\b &= 18.201(2) \text{ \AA} \\c &= 11.521(2) \text{ \AA} \\V &= 2434.6(5) \text{ \AA}^3\end{aligned}$$

For $Z = 4$ and F.W. = 462.50, the calculated density is 1.26 g/cm³. The systematic absences of:

$$\begin{aligned}h00: h &\neq 2n \\0k0: k &\neq 2n \\00l: l &\neq 2n\end{aligned}$$

uniquely determine the space group to be:

$$P2_12_12_1 \text{ (\#19)}$$

The data were collected at a temperature of $21 \pm 1^\circ\text{C}$ using the ω - 2θ scan technique to a maximum 2θ value of 120.1° . Omega scans of several intense reflections, made prior to data collection, had an average width at half-height of 0.22° with a take-off angle of 6.0° . Scans of $(1.52 + 0.30 \tan \theta)^\circ$ were made at a speed of $8.0^\circ/\text{min}$ (in omega). The weak reflections ($I < 10.0\sigma(I)$) were rescanned (maximum of 5 scans) and the counts were accumulated to ensure good counting statistics. Stationary background counts were recorded on each side of the reflection. The ratio of peak counting time to background counting time was 2:1. The diameter of the incident beam collimator was 0.5 mm and the crystal to detector distance was 235 mm. The computer-controlled slits were set to 3.0 mm (horizontal) and 3.0 mm (vertical).

Data Reduction

A total of 2094 reflections was collected. The intensities of three representative reflection were measured after every 150 reflections. No decay correction was applied.

The linear absorption coefficient, μ , for Cu-K α radiation is 8.0 cm^{-1} . Azimuthal scans of several reflections indicated no need for an absorption correction. The data were corrected for Lorentz and polarization effects.

Structure Solution and Refinement

The structure was solved by direct methods¹ and expanded using Fourier techniques². The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included but not refined. The final cycle of full-matrix least-squares refinement³ was based on 1926 observed reflections ($I > 2.00\sigma(I)$) and 298 variable parameters and converged (largest parameter shift was 0.09 times its esd) with unweighted and weighted agreement factors of:

$$R = \Sigma ||Fo| - |Fc|| / \Sigma |Fo| = 0.051$$

$$R_w = \sqrt{(\Sigma w(|Fo| - |Fc|)^2 / \Sigma w Fo^2)} = 0.068$$

The standard deviation of an observation of unit weight⁴ was 1.82. The weighting scheme was based on counting statistics and included a factor ($p = 0.060$) to downweight the intense reflections. Plots of $\Sigma w(|Fo| - |Fc|)^2$ versus $|Fo|$, reflection order in data collection, $\sin \theta/\lambda$ and various classes of indices showed no unusual trends. The maximum and minimum peaks on the final difference Fourier map corresponded to 0.21 and -0.32 $e^-/\text{\AA}^3$, respectively.

Neutral atom scattering factors were taken from Cromer and Waber⁵. Anomalous dispersion effects were included in F_{calc} ⁶; the values for $\Delta f'$ and $\Delta f''$ were those of Creagh and McAuley⁷. The values for the mass attenuation coefficients are those of Creagh and Hubbel⁸. All calculations were performed using the teXsan⁹ crystallographic software package of Molecular Structure Corporation.

References

- (1) SIR92: Altomare, A., Burla, M.C., Camalli, M., Cascarano, M., Giacovazzo, C., Guagliardi, A., Polidori, G. (1994). J. Appl. Cryst., ~~in preparation~~
27, P435 (1994)
- (2) DIRDIF94: Beurskens, P.T., Admiraal, G., Beurskens, G., Bosman, W.P., de Gelder, R., Israel, R. and Smits, J.M.M. (1994). The DIRDIF-94 program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.

- (3) Least-Squares:

$$\text{Function minimized: } \Sigma w(|Fo| - |Fc|)^2$$

$$\text{where } w = \frac{1}{\sigma_c^2(Fo)} = [\sigma_c^2(Fo) + \frac{p^2}{4} Fo^2]^{-1}$$

$$\sigma_c(Fo) = \text{e.s.d. based on counting statistics}$$

$$p = \text{p-factor}$$

- (4) Standard deviation of an observation of unit weight:

$$\sqrt{\Sigma w(|Fo| - |Fc|)^2 / (No - Nv)}$$

$$\text{where: } No = \text{number of observations}$$

$$Nv = \text{number of variables}$$

(5) Cromer, D. T. & Waber, J. T.; "International Tables for X-ray Crystallography", Vol. IV, The Kynoch Press, Birmingham, England, Table 2.2 A (1974).

(6) Ibers, J. A. & Hamilton, W. C.; *Acta Crystallogr.*, 17, 781 (1964).

(7) Creagh, D. C. & McAuley, W.J. ; "International Tables for Crystallography", Vol C, (A.J.C. Wilson, ed.), Kluwer Academic Publishers, Boston, Table 4.2.6.8, pages 219-222 (1992).

(8) Creagh, D. C. & Hubbell, J.H.; "International Tables for Crystallography", Vol C, (A.J.C. Wilson, ed.), Kluwer Academic Publishers, Boston, Table 4.2.4.3, pages 200-206 (1992).

(9) teXsan: Crystal Structure Analysis Package, Molecular Structure Corporation (1985 & 1992).

EXPERIMENTAL DETAILS

A. Crystal Data

Empirical Formula	$C_{23}H_{30}N_2O_8$
Formula Weight	462.50
Crystal Color, Habit	colorless, prism
Crystal Dimensions	0.35 X 0.20 X 0.20 mm
Crystal System	orthorhombic
Lattice Type	Primitive
No. of Reflections Used for Unit Cell Determination (2θ range)	25 (57.3 - 59.9°)
Omega Scan Peak Width at Half-height	0.22°
Lattice Parameters	$a = 11.610(2) \text{ \AA}$ $b = 18.201(2) \text{ \AA}$ $c = 11.521(2) \text{ \AA}$
	$V = 2434.6(5) \text{ \AA}^3$
Space Group	$P2_12_12_1$ (#19)
Z value	4
D_{calc}	1.262 g/cm ³
F_{000}	984.00
$\mu(\text{CuK}\alpha)$	8.02 cm ⁻¹

B. Intensity Measurements

Diffractometer	Rigaku AFC7R
Radiation	$\text{CuK}\alpha$ ($\lambda = 1.54178 \text{ \AA}$) graphite monochromated
Attenuator	Ni foil (factor = 9.38)

Take-off Angle	6.0°
Detector Aperture	3.0 mm horizontal 3.0 mm vertical
Crystal to Detector Distance	235 mm
Voltage, Current	40kV, 80mA
Temperature	21.0°C
Scan Type	ω - 2θ
Scan Rate	8.0°/min (in ω) (up to 5 scans)
Scan Width	$(1.52 + 0.30 \tan \theta)^\circ$
$2\theta_{max}$	120.1°
No. of Reflections Measured	Total: 2094
Corrections	Lorentz-polarization

C. Structure Solution and Refinement

Structure Solution	Direct Methods (SIR92)
Refinement	Full-matrix least-squares
Function Minimized	$\Sigma w(Fo - Fc)^2$
Least Squares Weights	$w = \frac{1}{\sigma^2(F_o)} = [\sigma_c^2(F_o) + \frac{p^2}{4} F_o^2]^{-1}$
p-factor	0.0600
Anomalous Dispersion	All non-hydrogen atoms
No. Observations ($I > 2.00\sigma(I)$)	1926
No. Variables	298
Reflection/Parameter Ratio	6.46
Residuals: R; Rw	0.051 ; 0.068
Goodness of Fit Indicator	1.82
Max Shift/Error in Final Cycle	0.09
Maximum peak in Final Diff. Map	0.21 $e^-/\text{\AA}^3$

Minimum peak in Final Diff. Map

-0.32 $e^-/\text{\AA}^3$

Table 1. Atomic coordinates and B_{iso}/B_{eq}

atom	x	y	z	B_{eq}
O(1)	0.3433(2)	-0.0984(1)	1.0692(2)	4.08(5)
O(2)	0.1277(2)	-0.1685(1)	1.0334(2)	4.18(5)
O(3)	0.0674(3)	0.1171(2)	0.8467(3)	5.43(7)
O(4)	0.2221(2)	0.1332(1)	0.9661(2)	4.36(6)
O(5)	0.4917(2)	0.0745(2)	1.0420(2)	4.54(6)
O(6)	0.3889(3)	-0.0536(2)	0.7175(3)	6.08(7)
O(7)	0.4931(3)	-0.1672(2)	0.9907(4)	7.15(9)
O(8)	0.0131(3)	-0.1385(2)	1.1943(2)	5.83(7)
N(1)	0.2003(2)	0.0272(2)	0.8736(3)	3.53(6)
N(2)	0.4090(2)	0.0102(1)	0.8903(2)	3.25(6)
C(1)	0.1550(3)	0.0955(2)	0.8925(3)	3.87(8)
C(2)	0.2949(3)	-0.0017(2)	0.9424(3)	3.03(6)
C(3)	0.2765(3)	-0.0830(2)	0.9684(3)	3.32(6)
C(4)	0.1475(3)	-0.0965(2)	0.9889(3)	3.44(7)
C(5)	0.0884(3)	-0.0874(2)	0.8742(3)	3.74(7)
C(6)	0.1347(3)	-0.0248(2)	0.8021(3)	3.71(7)
C(7)	0.0119(5)	-0.1346(3)	0.8331(4)	5.8(1)
C(8)	0.1913(4)	0.2087(2)	1.0024(4)	4.87(9)
C(9)	0.2820(5)	0.2238(3)	1.0938(5)	6.2(1)
C(10)	0.2008(7)	0.2592(3)	0.9005(5)	8.2(2)
C(11)	0.0723(5)	0.2101(3)	1.0569(5)	7.0(1)
C(12)	0.4995(3)	0.0440(2)	0.9509(3)	3.41(7)
C(13)	0.6041(3)	0.0312(2)	0.8798(3)	3.73(7)
C(14)	0.7172(3)	0.0527(2)	0.8987(4)	4.65(9)

Table 1. Atomic coordinates and B_{iso}/B_{eq} (continued)

atom	x	y	z	B_{eq}
C(15)	0.7991(3)	0.0307(3)	0.8175(4)	5.9(1)
C(16)	0.7680(4)	-0.0115(4)	0.7229(5)	6.9(1)
C(17)	0.6539(4)	-0.0309(3)	0.7030(4)	6.4(1)
C(18)	0.5742(3)	-0.0085(2)	0.7832(4)	4.44(8)
C(19)	0.4475(3)	-0.0214(2)	0.7864(4)	4.25(8)
C(20)	0.4012(4)	-0.1665(2)	1.0631(4)	5.7(1)
C(21)	0.5967(6)	-0.1426(4)	1.0402(9)	9.6(2)
C(22)	0.1117(4)	-0.1715(3)	1.1555(4)	5.33(10)
C(23)	-0.0887(5)	-0.1711(4)	1.1542(5)	7.4(1)
H(1)	0.2939	0.0234	1.0143	2.9989
H(2)	0.1184	-0.0606	1.0421	3.3706
H(3)	0.3036	-0.1125	0.9053	3.3054
H(4)	0.1087	-0.2226	1.1796	5.6940
H(5)	0.1758	-0.1494	1.1937	5.6940
H(6)	0.4280	-0.1798	1.1407	6.3200
H(7)	0.3475	-0.2042	1.0404	6.3200
H(8)	0.7399	0.0822	0.9631	4.6498
H(9)	0.8763	0.0436	0.8229	6.5916
H(10)	0.6331	-0.0593	0.6362	6.7244
H(11)	0.8256	-0.0276	0.6680	6.8659
H(12)	-0.1535	-0.1460	1.1844	8.1395
H(13)	-0.0896	-0.1727	1.0739	8.1395
H(14)	-0.0913	-0.2215	1.1844	8.1395
H(15)	0.6210	-0.1736	1.1039	10.8211

Table 1. Atomic coordinates and B_{iso}/B_{eq} (continued)

atom	x	y	z	B_{eq}
H(16)	0.6595	-0.1423	0.9848	10.8211
H(17)	0.5898	-0.0935	1.0701	10.8211
H(18)	0.2761	0.1898	1.1557	6.5088
H(19)	0.3570	0.2225	1.0618	6.5088
H(20)	0.2702	0.2729	1.1261	6.5088
H(21)	0.0569	0.2599	1.0833	7.6312
H(22)	0.0177	0.1952	1.0050	7.6312
H(23)	0.0733	0.1795	1.1249	7.6312
H(24)	0.2766	0.2580	0.8659	8.6318
H(25)	0.1480	0.2427	0.8384	8.6318
H(26)	0.1816	0.3083	0.9175	8.6318
H(27)	0.0725	-0.0004	0.7622	3.8122
H(28)	0.1845	-0.0447	0.7399	3.8122
H(29)	-0.0104	-0.1778	0.8771	5.8150
H(30)	-0.0237	-0.1285	0.7583	5.8150

$$B_{eq} = \frac{8}{3}\pi^2(U_{11}(aa^*)^2 + U_{22}(bb^*)^2 + U_{33}(cc^*)^2 + 2U_{12}aa^*bb^* \cos \gamma + 2U_{13}aa^*cc^* \cos \beta + 2U_{23}bb^*cc^* \cos \alpha)$$

Table 2. Anisotropic Displacement Parameters

atom	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
O(1)	0.055(1)	0.045(1)	0.055(1)	0.009(1)	-0.010(1)	0.000(1)
O(2)	0.064(2)	0.047(1)	0.047(1)	-0.009(1)	0.006(1)	0.004(1)
O(3)	0.058(2)	0.059(1)	0.090(2)	0.014(1)	-0.022(2)	0.001(2)
O(4)	0.048(1)	0.040(1)	0.078(2)	0.004(1)	-0.006(1)	-0.010(1)
O(5)	0.052(1)	0.070(2)	0.050(1)	-0.008(1)	-0.001(1)	-0.019(1)
O(6)	0.061(2)	0.112(2)	0.058(2)	-0.021(2)	0.006(1)	-0.040(2)
O(7)	0.067(2)	0.094(2)	0.111(3)	0.022(2)	-0.008(2)	-0.020(2)
O(8)	0.076(2)	0.090(2)	0.056(2)	-0.019(2)	0.018(2)	-0.014(2)
N(1)	0.040(1)	0.041(1)	0.053(2)	-0.002(1)	-0.005(1)	-0.001(1)
N(2)	0.039(1)	0.047(1)	0.038(1)	-0.005(1)	0.002(1)	-0.007(1)
C(1)	0.044(2)	0.041(2)	0.062(2)	0.001(2)	-0.002(2)	0.005(2)
C(2)	0.039(1)	0.039(2)	0.037(1)	0.002(1)	-0.001(1)	-0.003(1)
C(3)	0.042(2)	0.040(2)	0.044(2)	0.004(1)	0.002(2)	-0.006(1)
C(4)	0.047(2)	0.041(2)	0.042(2)	-0.004(1)	0.003(1)	-0.003(1)
C(5)	0.049(2)	0.047(2)	0.046(2)	-0.006(2)	-0.003(2)	-0.002(1)
C(6)	0.048(2)	0.047(2)	0.046(2)	-0.005(1)	-0.008(2)	0.001(2)
C(7)	0.085(3)	0.076(3)	0.061(2)	-0.036(3)	-0.018(2)	0.010(2)
C(8)	0.074(3)	0.038(2)	0.073(3)	0.009(2)	0.004(2)	-0.005(2)
C(9)	0.085(3)	0.061(2)	0.091(3)	0.001(2)	-0.005(3)	-0.021(2)
C(10)	0.187(7)	0.049(2)	0.075(3)	0.006(3)	0.009(5)	-0.001(2)
C(11)	0.073(3)	0.088(3)	0.105(4)	0.027(3)	0.003(3)	-0.018(3)
C(12)	0.041(2)	0.046(2)	0.042(2)	-0.001(1)	-0.003(2)	0.000(2)
C(13)	0.040(2)	0.051(2)	0.051(2)	0.001(2)	0.001(2)	0.002(2)
C(14)	0.042(2)	0.073(2)	0.061(2)	-0.007(2)	-0.002(2)	0.000(2)

Table 2. Anisotropic Displacement Parameters (continued)

atom	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
C(15)	0.038(2)	0.101(3)	0.083(3)	-0.004(2)	0.009(2)	-0.001(3)
C(16)	0.050(2)	0.126(4)	0.086(3)	-0.001(3)	0.022(2)	-0.021(3)
C(17)	0.063(3)	0.116(4)	0.066(3)	-0.010(3)	0.022(2)	-0.032(3)
C(18)	0.047(2)	0.066(2)	0.056(2)	-0.005(2)	0.007(2)	-0.012(2)
C(19)	0.048(2)	0.064(2)	0.050(2)	-0.008(2)	0.008(2)	-0.014(2)
C(20)	0.075(3)	0.057(2)	0.086(3)	0.025(2)	-0.007(3)	0.009(2)
C(21)	0.086(4)	0.097(4)	0.182(7)	-0.009(3)	-0.049(5)	0.017(5)
C(22)	0.080(3)	0.073(2)	0.050(2)	-0.015(2)	0.000(2)	0.011(2)
C(23)	0.082(3)	0.120(4)	0.077(3)	-0.028(3)	0.026(3)	-0.026(3)

The general temperature factor expression:

$$\exp(-2\pi^2(a^2U_{11}h^2 + b^2U_{22}k^2 + c^2U_{33}l^2 + 2a^*b^*U_{12}hk + 2a^*c^*U_{13}hl + 2b^*c^*U_{23}kl))$$

Table 3. Bond Lengths(Å)

atom	atom	distance	atom	atom	distance
O(1)	C(3)	1.425(4)	O(1)	C(20)	1.411(5)
O(2)	C(4)	1.426(4)	O(2)	C(22)	1.421(5)
O(3)	C(1)	1.211(5)	O(4)	C(1)	1.340(4)
O(4)	C(8)	1.481(4)	O(5)	C(12)	1.191(4)
O(6)	C(19)	1.199(5)	O(7)	C(20)	1.355(6)
O(7)	C(21)	1.404(7)	O(8)	C(22)	1.368(6)
O(8)	C(23)	1.402(6)	N(1)	C(1)	1.368(5)
N(1)	C(2)	1.453(4)	N(1)	C(6)	1.468(4)
N(2)	C(2)	1.470(4)	N(2)	C(12)	1.403(4)
N(2)	C(19)	1.402(5)	C(2)	C(3)	1.525(4)
C(3)	C(4)	1.536(5)	C(4)	C(5)	1.498(5)
C(5)	C(6)	1.509(5)	C(5)	C(7)	1.324(6)
C(8)	C(9)	1.515(7)	C(8)	C(10)	1.494(7)
C(8)	C(11)	1.517(7)	C(12)	C(13)	1.482(5)
C(13)	C(14)	1.388(5)	C(13)	C(18)	1.371(6)
C(14)	C(15)	1.393(6)	C(15)	C(16)	1.381(8)
C(16)	C(17)	1.390(7)	C(17)	C(18)	1.370(6)
C(18)	C(19)	1.489(5)			

Table 4. Bond Lengths(Å)

atom	atom	distance	atom	atom	distance
C(2)	H(1)	0.94	C(3)	H(3)	0.96
C(4)	H(2)	0.96	C(6)	H(27)	0.96
C(6)	H(28)	0.99	C(7)	H(29)	0.97
C(7)	H(30)	0.96	C(9)	H(18)	0.95
C(9)	H(19)	0.95	C(9)	H(20)	0.98
C(10)	H(24)	0.97	C(10)	H(25)	0.99
C(10)	H(26)	0.94	C(11)	H(21)	0.97
C(11)	H(22)	0.91	C(11)	H(23)	0.96
C(14)	H(8)	0.95	C(15)	H(9)	0.93
C(16)	H(11)	0.97	C(17)	H(10)	0.96
C(20)	H(6)	0.98	C(20)	H(7)	0.96
C(21)	H(15)	0.97	C(21)	H(16)	0.97
C(21)	H(17)	0.96	C(22)	H(4)	0.97
C(22)	H(5)	0.95	C(23)	H(12)	0.95
C(23)	H(13)	0.93	C(23)	H(14)	0.98

Table 5. Bond Angles(°)

atom	atom	atom	angle	atom	atom	atom	angle
C(3)	O(1)	C(20)	113.0(3)	C(4)	O(2)	C(22)	114.4(3)
C(1)	O(4)	C(8)	120.9(3)	C(20)	O(7)	C(21)	115.0(5)
C(22)	O(8)	C(23)	114.3(3)	C(1)	N(1)	C(2)	122.2(3)
C(1)	N(1)	C(6)	118.4(3)	C(2)	N(1)	C(6)	117.8(3)
C(2)	N(2)	C(12)	122.5(3)	C(2)	N(2)	C(19)	125.1(3)
C(12)	N(2)	C(19)	111.4(3)	O(3)	C(1)	O(4)	126.7(3)
O(3)	C(1)	N(1)	123.3(4)	O(4)	C(1)	N(1)	110.0(3)
N(1)	C(2)	N(2)	113.9(3)	N(1)	C(2)	C(3)	110.6(3)
N(2)	C(2)	C(3)	110.4(3)	O(1)	C(3)	C(2)	105.9(3)
O(1)	C(3)	C(4)	111.9(3)	C(2)	C(3)	C(4)	108.8(3)
O(2)	C(4)	C(3)	111.1(3)	O(2)	C(4)	C(5)	110.2(3)
C(3)	C(4)	C(5)	107.0(3)	C(4)	C(5)	C(6)	114.0(3)
C(4)	C(5)	C(7)	123.4(3)	C(6)	C(5)	C(7)	122.1(3)
N(1)	C(6)	C(5)	111.2(3)	O(4)	C(8)	C(9)	101.4(3)
O(4)	C(8)	C(10)	109.3(4)	O(4)	C(8)	C(11)	110.6(4)
C(9)	C(8)	C(10)	112.5(4)	C(9)	C(8)	C(11)	110.0(4)
C(10)	C(8)	C(11)	112.5(5)	O(5)	C(12)	N(2)	125.8(3)
O(5)	C(12)	C(13)	128.5(3)	N(2)	C(12)	C(13)	105.7(3)
C(12)	C(13)	C(14)	130.1(4)	C(12)	C(13)	C(18)	108.9(3)
C(14)	C(13)	C(18)	121.0(4)	C(13)	C(14)	C(15)	117.4(4)
C(14)	C(15)	C(16)	120.7(4)	C(15)	C(16)	C(17)	121.4(4)
C(16)	C(17)	C(18)	117.2(4)	C(13)	C(18)	C(17)	122.2(4)
C(13)	C(18)	C(19)	108.2(3)	C(17)	C(18)	C(19)	129.5(4)
O(6)	C(19)	N(2)	125.8(3)	O(6)	C(19)	C(18)	128.4(4)

Table 5. Bond Angles(°) (continued)

atom	atom	atom	angle	atom	atom	atom	angle
N(2)	C(19)	C(18)	105.7(3)	O(1)	C(20)	O(7)	114.5(4)
O(2)	C(22)	O(8)	114.6(4)				

Table 6. Bond Angles(°)

atom	atom	atom	angle	atom	atom	atom	angle
N(1)	C(2)	H(1)	107.1	N(2)	C(2)	H(1)	107.3
C(3)	C(2)	H(1)	107.1	O(1)	C(3)	H(3)	109.3
C(2)	C(3)	H(3)	110.5	C(4)	C(3)	H(3)	110.4
O(2)	C(4)	H(2)	109.9	C(3)	C(4)	H(2)	109.5
C(5)	C(4)	H(2)	109.1	N(1)	C(6)	H(27)	111.1
N(1)	C(6)	H(28)	109.8	C(5)	C(6)	H(27)	110.1
C(5)	C(6)	H(28)	109.3	H(27)	C(6)	H(28)	105.1
C(5)	C(7)	H(29)	121.2	C(5)	C(7)	H(30)	122.2
H(29)	C(7)	H(30)	116.6	C(8)	C(9)	H(18)	110.8
C(8)	C(9)	H(19)	111.4	C(8)	C(9)	H(20)	109.4
H(18)	C(9)	H(19)	110.1	H(18)	C(9)	H(20)	107.5
H(19)	C(9)	H(20)	107.5	C(8)	C(10)	H(24)	112.2
C(8)	C(10)	H(25)	109.6	C(8)	C(10)	H(26)	113.7
H(24)	C(10)	H(25)	105.1	H(24)	C(10)	H(26)	108.8
H(25)	C(10)	H(26)	107.0	C(8)	C(11)	H(21)	108.2
C(8)	C(11)	H(22)	110.9	C(8)	C(11)	H(23)	108.5
H(21)	C(11)	H(22)	110.7	H(21)	C(11)	H(23)	106.6
H(22)	C(11)	H(23)	111.8	C(13)	C(14)	H(8)	122.8
C(15)	C(14)	H(8)	119.8	C(14)	C(15)	H(9)	122.8
C(16)	C(15)	H(9)	116.5	C(15)	C(16)	H(11)	120.2
C(17)	C(16)	H(11)	118.3	C(16)	C(17)	H(10)	120.7
C(18)	C(17)	H(10)	122.2	O(1)	C(20)	H(6)	108.8
O(1)	C(20)	H(7)	109.4	O(7)	C(20)	H(6)	108.0
O(7)	C(20)	H(7)	109.6	H(6)	C(20)	H(7)	106.1

Table 6. Bond Angles(°) (continued)

atom	atom	atom	angle	atom	atom	atom	angle
O(7)	C(21)	H(15)	111.8	O(7)	C(21)	H(16)	112.3
O(7)	C(21)	H(17)	111.8	H(15)	C(21)	H(16)	106.5
H(15)	C(21)	H(17)	107.1	H(16)	C(21)	H(17)	107.0
O(2)	C(22)	H(4)	109.0	O(2)	C(22)	H(5)	109.7
O(8)	C(22)	H(4)	107.3	O(8)	C(22)	H(5)	108.4
H(4)	C(22)	H(5)	107.5	O(8)	C(23)	H(12)	110.2
O(8)	C(23)	H(13)	110.6	O(8)	C(23)	H(14)	107.8
H(12)	C(23)	H(13)	112.0	H(12)	C(23)	H(14)	107.3
H(13)	C(23)	H(14)	108.9				

Table 7. Non-bonded Contacts out to 3.60 Å

atom	atom	distance	ADC	atom	atom	distance	ADC
O(1)	O(3)	3.378(4)	2	O(1)	C(6)	3.506(4)	2
O(2)	O(7)	3.384(4)	44703	O(2)	C(21)	3.558(7)	44703
O(3)	C(20)	3.409(6)	55402	O(3)	C(15)	3.505(5)	45501
O(4)	O(6)	3.485(4)	2	O(5)	C(6)	3.457(4)	2
O(5)	C(7)	3.528(5)	2	O(6)	C(11)	3.426(7)	55402
O(7)	C(23)	3.514(7)	54703	O(8)	C(18)	3.039(5)	2
O(8)	C(19)	3.130(5)	2	O(8)	C(13)	3.199(5)	2
O(8)	N(2)	3.372(4)	2	O(8)	C(12)	3.423(4)	2
C(18)	C(23)	3.596(7)	55402				

The ADC (atom designator code) specifies the position of an atom in a crystal. The 5-digit number shown in the table is a composite of three one-digit numbers and one two-digit number: TA (first digit) + TB (second digit) + TC (third digit) + SN (last two digits). TA, TB and TC are the crystal lattice translation digits along cell edges a, b and c. A translation digit of 5 indicates the origin unit cell. If TA = 4, this indicates a translation of one unit cell length along the a-axis in the negative direction. Each translation digit can range in value from 1 to 9 and thus ± 4 lattice translations from the origin (TA=5, TB=5, TC=5) can be represented.

The SN, or symmetry operator number, refers to the number of the symmetry operator used to generate the coordinates of the target atom. A list of symmetry operators relevant to this structure are given below.

For a given intermolecular contact, the first atom (origin atom) is located in the origin unit cell and its position can be generated using the identity operator (SN=1). Thus, the ADC for an origin atom is always 55501. The position of the second atom (target atom) can be generated using the ADC and the coordinates of the atom in the parameter table. For example, an ADC of 47502 refers to the target atom moved through symmetry operator two, then translated -1 cell translations along the a axis, +2 cell translations along the b axis, and 0 cell translations along the c axis.

An ADC of 1 indicates an intermolecular contact between two fragments (eg. cation and anion) that reside in the same asymmetric unit.

Symmetry Operators:

(1)	X,	Y,	Z	(2)	1/2-X,	-Y,	1/2+Z
(3)	1/2+X,	1/2-Y,	-Z	(4)	-X,	1/2+Y,	1/2-Z