Mechanism Underlying the Onset of Internal Blue Discoloration in Japanese Radish (*Raphanus sativus*) Roots

Katsunori Teranishi^{*†}, Nagata Masayasu[‡], and Daisuke Masuda[§]

[†]Graduate School of Bioresources, Mie University, 1577 Kurimamachiya, Tsu, Mie, Japan
[‡]Food Research Institute, National Agriculture and Food Research Organization, 2-1-12 Kannondai,
Tsukuba, Ibaraki, Japan

[§]Sand Dune Agricultural Research Center, Agricultural Experiment Station, Ishikawa Agriculture and Forestry Research Center, I 5-2 Uchihisumi, Kahoku, Ishikawa, Japan

Corresponding author Address:

Graduate School of Bioresources, Mie University, 1577 Kurimamachiya, Tsu, Mie, Japan. Tel: +81-59-231-9615; fax: +81-59-231-9615. E-mail address: teranisi@bio.mie-u.ac.jp.



Figure S1. Three groups of blue discoloration from 18 cultivars of Japanese radish roots. Upper photos for each cultivar: after storage at 20 °C for 4 day. Bottom photos for each

cultivar: after soaking in 0.29 M aqueous H_2O_2 solution at room temperature for 10 min. Left group is for cultivars that showed the discoloration with both treatments. Center group is for cultivars that showed no discoloration with storage at 20 °C and the discoloration with the H_2O_2 treatment. Right group is for cultivars that showed little or no blue discoloration with both treatments.