

Supplementary Information for

Incorporation of technetium into spinel ferrites

Wayne W. Lukens,^{1*} Nicola Magnani,^{1,2} Tolek Tyliczszak,³ Carolyn I. Pearce,⁴ David K. Shuh¹

- 1) Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720
- 2) European Commission, Joint Research Centre, Institute for Transuranium Elements, Karlsruhe
- 3) Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA 94720
- 4) Geosciences Group, Pacific Northwest National Laboratory, 902 Battelle Boulevard, K6-81, Richland, WA 99354

* MS 70A-1150, Lawrence Berkeley National Laboratory, Berkeley, CA 94720. Phone (510) 486-4305; Fax (510) 486-5596. WWLukens@lbl.gov.

Contents of SI:

Table S1. Analytical data for Tc-doped spinel ferrites	S2
Table S2. K_d values for the synthesis of Tc-doped spinel ferrites	S3
Spectra and data for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$	S4
Spectra and data for $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$	S6
Spectra and data for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$	S8
Spectra and data for $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-o}$	S10
Spectra and data for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-o}$	S12
Spectra and data for $\text{Tc}_{0.1}\text{Fe}_{2.9}\text{O}_4\text{-o}$	S14
Spectra and data for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-o}$	S16
Spectra and data for $\text{Tc}_{0.1}\text{Ni}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-o}$	S18
Spectra and data for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$	S20
Spectra and data for $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$	S22
Spectra and data for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$	S24
Spectra and data for $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$	S26
Spectra and data for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$	S28
Spectra and data for $\text{Tc}_{0.1}\text{Fe}_{2.9}\text{O}_4\text{-c}$	S30
Spectra and data for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$	S32
Spectra and data for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{2.7}\text{O}_4\text{-c}$	S34
References	S36

Experimental

Table S1. Analytical data for Tc-doped spinel ferrites^a

Sample ^b	% TcO ₄ ⁻ removed ^c	%Tc recovered ^d	Size (nm) ^e	a (Å)	a (Å) (w/o Tc)	Fe(II)/Fe ^{tot}	Fe(II)/Fe ^{tot} (w/o Tc)
Tc _{0.1} Mn _{1.1} Fe _{1.8} O ₄ -o	99.975(2)	99(2)	19	8.4873(8)	8.4702(3)	0.11(2)	0.08(1)
Tc _{0.1} Co _{1.1} Fe _{1.8} O ₄ -o	99.921(3)	101(4)	42	8.4239(1)	8.4186(4)	0.18(5)	0.20(1)
Tc _{0.1} Ni _{1.1} Fe _{1.8} O ₄ -o	99.975(2)	97(1)	28	8.3881(4)	8.382(2)	0.05(1)	0.12(1)
Tc _{0.1} Mg _{0.2} Fe _{2.7} O ₄ -o	99.984(2)	111(4)	38	8.4017(1)	8.4009(5)	0.22(2)	0.25(1)
Tc _{0.1} Mn _{0.2} Fe _{2.7} O ₄ -o	99.909(4)	94(1)	41	8.4205(1)	8.4160(5)	0.22(1)	0.20(1)
Tc _{0.1} Fe _{2.9} O ₄ -o	99.937(3)	89(1)	42	8.3916(1)	8.3987(6)	0.22(1)	0.26(1)
Tc _{0.1} Co _{0.2} Fe _{2.7} O ₄ -o	99.884(4)	92(1)	46	8.4010(1)	8.3993(7)	0.27(1)	0.26(1)
Tc _{0.1} Ni _{0.2} Fe _{2.7} O ₄ -o	99.901(4)	94(1)	36	8.3936(1)	8.3927(5)	0.24(1)	0.22(1)
Tc _{0.1} Mn _{1.1} Fe _{1.8} O ₄ -c	99.763(7)	81(1)	24	8.4906(4)	8.481(1)	0.02(1)	0.09(1)
Tc _{0.1} Co _{1.1} Fe _{1.8} O ₄ -c	99.49(1)	83(2)	9.2	8.4081(7)	8.415(1)	0.26(1)	0.37(1)
Tc _{0.1} Ni _{1.1} Fe _{1.8} O ₄ -c	99.52(1)	93(2)	7.7	8.374(3)	8.372(1)	0.10(1)	0.13(1)
Tc _{0.1} Mg _{0.2} Fe _{2.7} O ₄ -c	99.884(5)	96(2)	9.2	8.375(1)	8.3966(6)	0.22(2)	0.34(1)
Tc _{0.1} Mn _{0.2} Fe _{2.7} O ₄ -c	99.833(6)	100(2)	13	8.3932(7)	8.4143(3)	0.18(2)	0.32(1)
Tc _{0.1} Fe _{2.9} O ₄ -c	99.820(6)	88(2)	12	8.3672(3)	8.3946(4)	0.23(1)	0.37(1)
Tc _{0.1} Co _{0.2} Fe _{2.7} O ₄ -c	99.750(7)	101(1)	10	8.3767(8)	8.3962(4)	0.25(1)	0.36(1)
Tc _{0.1} Ni _{0.2} Fe _{2.7} O ₄ -c	99.848(6)	95(2)	12	8.3916(8)	8.3869(4)	0.22(1)	0.31(1)

- a) Standard deviations are given in parentheses in the same units as the last digit
b) Samples with an “o” suffix were prepared by oxidation; “c” by coprecipitation
c) TcO₄⁻ removed from solution during formation of Tc-doped spinel ferrite
d) Tc recovered by dissolving Tc-doped spinel ferrite
e) Crystallite (particle) diameter determined by Rietveld analysis

Table S2. K_d values for the synthesis of Tc-doped spinel ferrites^a

Sample ^b	K_d (L/kg)
$Tc_{0.1}Mn_{1.1}Fe_{1.8}O_4-o$	1.07E+05
$Tc_{0.1}Co_{1.1}Fe_{1.8}O_4-o$	3.32E+04
$Tc_{0.1}Ni_{1.1}Fe_{1.8}O_4-o$	1.05E+05
$Tc_{0.1}Mg_{0.2}Fe_{2.7}O_4-o$	1.97E+05
$Tc_{0.1}Mn_{0.2}Fe_{2.7}O_4-o$	3.37E+04
$Tc_{0.1}Fe_{2.9}O_4-o$	5.00E+04
$Tc_{0.1}Co_{0.2}Fe_{2.7}O_4-o$	2.63E+04
$Tc_{0.1}Ni_{0.2}Fe_{2.7}O_4-o$	3.09E+04
$Tc_{0.1}Mn_{1.1}Fe_{1.8}O_4-c$	1.41E+04
$Tc_{0.1}Co_{1.1}Fe_{1.8}O_4-c$	6.41E+03
$Tc_{0.1}Ni_{1.1}Fe_{1.8}O_4-c$	6.82E+03
$Tc_{0.1}Mg_{0.2}Fe_{2.7}O_4-c$	3.32E+04
$Tc_{0.1}Mn_{0.2}Fe_{2.7}O_4-c$	2.24E+04
$Tc_{0.1}Fe_{2.9}O_4-c$	2.11E+04
$Tc_{0.1}Co_{0.2}Fe_{2.7}O_4-c$	1.49E+04
$Tc_{0.1}Ni_{0.2}Fe_{2.7}O_4-c$	2.49E+04

a) Note that K_d may not be particularly meaningful in this context since ^{99}Tc is incorporated into the lattice of the solid rather than being adsorbed on the surface.

b) Samples with an “o” suffix were prepared by oxidation; “c” by coprecipitation




Figure S1: XRD pattern for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_{4-\text{o}}$ (Data in color; Rietveld fit in black)





Figure S2: XANES fit for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_{4-\text{o}}$

Table S1: Best EXAFS fit parameters $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.28(8)	0.001	1.59(1)	--	<0.001
O	5.72(8) ^b	0.0038(2)	2.025(3)	2.06	<0.001
Fe	4.0(1)	0.0035(8)	3.087(7)	2.97	<0.001
Tc	2.0(1) ^b	0.0035(8) ^c	3.087(7) ^c	2.97	0.018
Fe	6	0.0125(9)	3.565(8)	3.48	<0.001
O	8	0.021(6)	3.26(3)	3.57	0.009
O-Tc-O (MS)	6	0.016(2)	4.21(3)	4.12	0.002
Fe	12	0.016(2) ^c	5.27(2)	5.14	0.001

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 4.0(5)$ eV, $\chi^2 = 828$, $\chi_r^2 = 53.7$, $R = 0.005$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S3:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$




Figure S4: XRD pattern for Tc_{0.1}Co_{1.1}Fe_{1.8}O₄-o (Data in color; Rietveld fit in black)




Figure S5: XANES fit for Tc_{0.1}Co_{1.1}Fe_{1.8}O₄-o


Table S2: Best EXAFS fit parameters $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.3(1)	0.001	1.54(3)	--	0.088
O ^c	5.7(1) ^b	0.0027(3)	2.009(4)	2.06	<0.001
Fe	6	0.0059(4)	3.035(5)	2.97	<0.001
Fe	6	0.0058(5)	3.516(7)	3.48	<0.001
O-Tc-O (MS)	6	0.008(1)	4.16(6)	4.12	0.141
O	12	0.008(1) ^c	4.73(3)	4.76	0.031
Fe	12	0.008(1) ^c	5.19(1)	5.14	<0.001

a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = -2.1(8)$ eV, $\chi^2 = 1875$, $\chi^2_{\nu} = 107.7$, $R = 0.018$.

b) Parameter is constrained to equal 6 minus the preceding parameter

c) Parameter constrained to equal the preceding parameter

**Figure S6:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-o}$





Figure S7: XRD pattern for Tc_{0.1}Ni_{1.1}Fe_{1.8}O_{4-o} (Data in color; Rietveld fit in black)




Figure S8: XANES fit for Tc_{0.1}Ni_{1.1}Fe_{1.8}O_{4-o}


Table S3: Best EXAFS fit parameters for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_{4-o}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.4(1)	0.001	1.79(3)	--	0.070
O	5.6(1) ^b	0.0026(4)	2.009(4)	2.06	<0.001
Fe	4.8(5)	0.0029(2)	3.034(6)	2.97	<0.001
Tc	1.2(5) ^b	0.0029(2) ^c	3.034(6) ^c	2.97	0.101
Fe	6	0.0066(5)	3.514(7)	3.48	<0.001
O-Tc-O (MS)	5.6(1)	0.008(1)	4.12(5)	4.12	0.039
Fe	12	0.008(1) ^c	5.19(1)	5.14	<0.001

a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 13 parameters, $S_0^2 = 0.8$, $\Delta E_0 = -2(1)$ eV, $\chi^2 = 814$, $\chi_r^2 = 46.7$, $R = 0.013$. Parameters without errors are fixed.

b) Parameter is constrained to equal 6 minus the preceding parameter

c) Parameter constrained to equal the preceding parameter

**Figure S9:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_{4-o}$





Figure S10: XRD pattern for Tc_{0.1}Mg_{0.2}Fe_{2.7}O_{4-o} (Data in color; Rietveld fit in black)




Figure S11: XANES fit for Tc_{0.1}Mg_{0.2}Fe_{2.7}O_{4-o}


Table S4: Best EXAFS fit parameters for $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	P
O	0.29(9)	0.001	1.54(2)	--	0.012
O ^c	5.71(9) ^b	0.0031(3)	2.034(4)	2.06	<0.001
Fe	4.1(2)	0.004(2)	3.096(9)	2.97	<0.001
Tc	1.9(2) ^b	0.004(2) ^c	3.096(9) ^b	2.97	0.077
Fe	6	0.0070(5)	3.537(7)	3.48	<0.001
O-Tc-O (MS)	6	0.011(1)	4.21(4)	4.12	0.003
O	12	0.011(1) ^c	4.78(4)	4.76	0.039
Fe	12	0.011(1) ^c	5.22(2)	5.14	0.001

a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 2.0(8)$ eV, $\chi^2 = 4049$, $\chi_r^2 = 247$, $R = 0.013$.

b) Parameter is constrained to equal 6 minus the preceding parameter

c) Parameter constrained to equal the preceding parameter

**Figure S12:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$





Figure S13: XRD pattern for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$ (Data in color; Rietveld fit in black)





Figure S14: XANES fit for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$

Table S5: Best EXAFS fit parameters for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	O_h Fe_3O_4 ($\text{\AA}$)	p
O	0.38(9)	0.001	1.51(1)	--	0.026
O	5.62(9) ^b	0.0043(3)	2.036(4)	2.06	<0.001
Fe	3.80(8)	0.0017(6)	3.100(8)	2.97	<0.001
Tc	2.20(8) ^b	0.0017(6) ^c	3.100(8) ^c	2.97	<0.001
Fe	6	0.0100(9)	3.53(1)	3.48	<0.001
O-Tc-O (MS)	6	0.013(2)	4.22(5)	4.12	0.003
Fe	12	0.013(2) ^c	5.25(2)	5.14	0.004

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 13 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 6.0(9)$ eV, $\chi^2 = 572$ $\chi_v^2 = 32.9$ $R = 0.020$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S15:** Tc K-edge EXAFS spectrum of Tc in for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$

Tc_{0.1}Fe_{2.9}O_{4-o}




Figure S16: XRD pattern for Tc_{0.1}Fe_{2.9}O_{4-o} (Data in color; Rietveld fit in black)





Figure S17: XANES fit for Tc_{0.1}Fe_{2.9}O_{4-o} (XANES data was not fit. The XANES spectrum of this sample is the Tc(IV) in spinel ferrite standard spectrum)

Table S6: Best EXAFS fit parameters for $\text{Tc}_{0.1}\text{Fe}_{2.9}\text{O}_{4-\text{o}}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.20(8)	0.001	1.61(2)	--	0.002
O	5.80(8) ^b	0.0039(2)	2.027(3)	2.06	<0.001
Fe	3.9(1)	0.0036(9)	3.090(7)	2.97	<0.001
Tc	2.1(1) ^b	0.0036(9) ^c	3.090(7) ^c	2.97	0.001
Fe	6	0.0084(5)	3.524(6)	3.48	<0.001
O	8	0.009(3)	4.07(2)	3.57	0.262
O-Tc-O (MS)	6	0.013(2)	4.27(5)	4.12	0.051
Fe	12	0.013(2) ^c	5.22(2)	5.14	<0.001

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 2.1(7)$ eV, $\chi^2 = 434$, $\chi^2_{\nu} = 28.4$, $R = 0.009$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S18:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Fe}_{2.9}\text{O}_{4-\text{o}}$





Figure S19: XRD pattern for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$ (Data in color; Rietveld fit in black)





Figure S20: XANES fit for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$

Table S7: Best fit parameters for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$ ^a

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.4(1)	0.001	1.66(2)	--	<0.001
O	5.6(1) ^b	0.0046(5)	2.026(6)	2.06	<0.001
Fe	3.8(1)	0.003(1)	3.08(1)	2.97	<0.001
Tc	2.2(1) ^b	0.003(1) ^c	3.08(1) ^c	2.97	0.020
Fe	6	0.0084(9)	3.51(1)	3.48	<0.001
O-Tc-O (MS)	6	0.012(2)	4.19(6)	4.12	0.050
O	12	0.012(2) ^c	4.77(6)	4.76	0.256
Fe	12	0.012(2) ^c	5.20(2)	5.14	0.009

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 14 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 7(1)$ eV, $\chi^2 = 935$, $\chi_v^2 = 57$, $R = 0.031$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S21:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$





Figure S22: XRD pattern for Tc_{0.1}Ni_{0.2}Fe_{2.7}O_{4-o} (Data in color; Rietveld fit in black)




Figure S23: XANES fit for Tc_{0.1}Ni_{0.2}Fe_{2.7}O_{4-o}


Table S8: Best fit parameters for $\text{Tc}_{0.1}\text{Ni}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$ ^a

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.31(8)	0.001	1.57(1)	--	<0.001
O ^c	5.69(8)	0.0032(2)	2.030(3)	2.06	<0.001
Fe	4.1(1)	0.003(1)	3.075(7)	2.97	<0.001
Tc	1.9(1)	0.003(1) ^b	3.075(7) ^b	2.97	0.002
Fe	6	0.0074(5)	3.530(6)	3.48	<0.001
O-Tc-O (MS)	6	0.011(1)	4.21(3)	4.12	<0.001
O	12	0.011(1) ^b	4.77(3)	4.76	0.023
Fe	12	0.011(1) ^b	5.21(1)	5.14	<0.001

a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 14 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 4.2(7)$ eV, $\chi^2 = 1689$, $\chi_r^2 = 103$, $R = 0.009$.

b) Parameter is constrained to equal 6 minus the preceding parameter

c) Parameter constrained to equal the preceding parameter

**Figure S24:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Ni}_{0.2}\text{Fe}_{2.7}\text{O}_{4-\text{o}}$




Figure S25: XRD pattern for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_{4-\text{c}}$ (Data in color; Rietveld fit in black)





Figure S26: XANES fit for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_{4-\text{c}}$

Table S9: Best fit parameters for $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.4(1)	0.001	1.72(1)	--	0.007
O	5.6(1) ^b	0.0043(3)	2.016(5)	2.06	<0.001
Fe	4.0(2)	0.003(1)	3.083(1)	2.97	0.057
Tc	2.0(2) ^b	0.003(1) ^c	3.083(1) ^c	2.97	0.108
Fe	6	0.015(3)	3.550(3)	3.48	0.039
O	8	0.017(7)	3.28(5)	3.48	0.091
O-Tc-O (MS)	6	0.018(4)	4.14(7)	4.12	0.096
Fe	12	0.018(4) ^c	5.26(4)	5.14	0.044

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 3(1)$ eV, $\chi^2 = 4425$, $\chi_v^2 = 287$, $R = 0.020$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S27:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Mn}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$




Figure S28: XRD pattern for $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$ (Data in color; Rietveld fit in black)





Figure S29: XANES fit for $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$

Table S10: Best fit parameters for $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}^{\text{a}}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.84(7)	0.001	1.718(5)	--	<0.001
O	5.16(7) ^b	0.0044(4)	2.027(5)	2.06	<0.001
Fe	3.93(8)	0.0008(5)	3.069(8)	2.97	<0.001
Tc	2.07(8) ^b	0.0008(5) ^c	3.069(8) ^c	2.97	<0.001
Fe	6	0.011(1)	3.54(1)	3.48	<0.001
O-Tc-O (MS)	6	0.012(2)	4.21(5)	4.12	<0.001
Fe	12	0.012(2) ^c	5.23(2)	5.14	<0.001

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 13 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 2(1)$ eV, $\chi^2 = 1777$ $\chi_v^2 = 102$ $R = 0.020$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S30:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Co}_{1.1}\text{Fe}_{1.8}\text{O}_4\text{-c}$





Figure S31: XRD pattern for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_{4-\text{c}}$ (Data in color; Rietveld fit in black)





Figure S32: XANES fit for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_{4-\text{c}}$

Table S11: Best fit parameters for $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_{4-c}$ ^a

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	1.2(1)	0.001	1.726(6)	--	<0.001
O	4.8(1) ^b	0.0056(6)	2.027(8)	2.06	<0.001
Fe	3.7(2)	0.005(1)	3.10(2)	2.97	<0.001
Tc	2.3(2) ^b	0.005(1) ^c	3.10(2) ^c	2.97	0.038
Fe	6	0.018(4)	3.50(3)	3.48	0.037
O-Tc-O (MS)	6	0.01(1)	4.18(1)	4.12	0.085

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 12 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 4(2)$ eV, $\chi^2 = 3866$ $\chi_v^2 = 210$ $R = 0.058$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S33:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Ni}_{1.1}\text{Fe}_{1.8}\text{O}_{4-c}$




Figure S34: XRD pattern for $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$ (Data in color; Rietveld fit in black)





Figure S35: XANES fit for $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$

Table S12: Best fit parameters for $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}^{\text{a}}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.33(8)	0.001	1.671(6)	--	0.021
O	5.67(9) ^b	0.0048(3)	2.027(8)	2.06	<0.001
Fe	3.73(8)	0.0032(7)	3.10(2)	2.97	<0.001
Tc	2.27(8) ^b	0.0032(7) ^c	3.10(2) ^c	2.97	0.002
Fe	6	0.0116(9)	3.51(3)	3.48	<0.001
O	8	0.04(3)	3.44(3)	3.48	0.336
O-Tc-O (MS)	6	0.015(2)	4.17(1)	4.12	0.002
Fe	12	0.015(2)	5.23(1)	5.14	0.012

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 2(1)$ eV, $\chi^2 = 1083$, $\chi_{\nu}^2 = 70$, $R = 0.013$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S36:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Mg}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$





Figure S37: XRD pattern for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$ (Data in color; Rietveld fit in black)




Figure S38: XANES fit for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$


Table S13: Best fit parameters for $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}^a$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.66(6)	0.001	1.715(7)	--	<0.001
O	5.34(6) ^b	0.0051(3)	2.019(4)	2.06	<0.001
Fe	3.90(8)	0.0031(7)	3.083(7)	2.97	<0.001
Tc	2.10(8) ^b	0.0031(7) ^c	3.083(7) ^c	2.97	<0.001
Fe	6	0.0117(8)	3.509(9)	3.48	<0.001
O-Tc-O (MS)	6	0.014(2)	4.17(4)	4.12	<0.001
Fe	12	0.014(2)	5.23(2)	5.14	0.002

a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 13 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 5.4(8)$ eV, $\chi^2 = 187$ $\chi_v^2 = 10.7$ $R = 0.013$. Parameters without errors are fixed.

b) Parameter is constrained to equal 6 minus the preceding parameter

c) Parameter constrained to equal the preceding parameter

**Figure S39:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Mn}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$

Tc_{0.1}Fe_{2.9}O₄-c




Figure S40: XRD pattern for Tc_{0.1}Fe_{2.9}O₄-c (Data in color; Rietveld fit in black)




Figure S41: XANES fit for Tc_{0.1}Fe_{2.9}O₄-c

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	O_h Fe_3O_4 ($\text{\AA}$)	p
O	0.29(6)	0.001	1.63(1)	--	0.001
O	5.71(6) ^b	0.0046(2)	2.020(3)	2.06	<0.001
Fe	3.83(6)	0.0024(6)	3.086(6)	2.97	<0.001
Tc	2.17(6) ^b	0.0024(6) ^c	3.086(6) ^c	2.97	<0.001
Fe	6	0.0113(7)	3.511(8)	3.48	<0.001
O-Tc-O (MS)	6	0.014(2)	4.17(3)	4.12	<0.001
Fe	12	0.014(2) ^c	5.23(2)	5.14	0.002

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 15 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 3(1)$ eV, $\chi^2 = 4425$ $\chi_r^2 = 287$ $R = 0.020$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter




Figure S42: Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Fe}_{2.9}\text{O}_4\text{-c}$




Figure S43: XRD pattern for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$ (Data in color; Rietveld fit in black)




Figure S44: XANES fit for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$


Table S15: Best fit parameters for $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}^{\text{a}}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.58(8)	0.001	1.716(9)	--	<0.001
O	5.42(8) ^b	0.0049(3)	2.020(5)	2.06	<0.001
Fe	3.84(9)	0.0029(8)	3.077(9)	2.97	<0.001
Tc	2.16(9) ^b	0.0029(8) ^c	3.077(9) ^c	2.97	0.001
Fe	6	0.0110(9)	3.51(1)	3.48	<0.001
O-Tc-O (MS)	6	0.013(2)	4.17(5)	4.12	0.003
Fe	12	0.013(2)	5.22(2)	5.14	0.003

a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 13 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 1(1)$ eV, $\chi^2 = 845$, $\chi_v^2 = 48.5$, $R = 0.021$. Parameters without errors are fixed.

b) Parameter is constrained to equal 6 minus the preceding parameter

c) Parameter constrained to equal the preceding parameter

**Figure S45:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Co}_{0.2}\text{Fe}_{2.7}\text{O}_4\text{-c}$





Figure S46: XRD pattern for Tc_{0.1}Ni_{0.2}Fe_{2.7}O₄-c (Data in color; Rietveld fit in black). Data was obtained using a Co source, but the 2θ values have been transformed to match the other data, which was obtained with a Cu source.





Figure S47: XANES fit for Tc_{0.1}Ni_{0.2}Fe_{2.7}O₄-c

Table S16: Best fit parameters $\text{Tc}_{0.1}\text{Ni}_{0.2}\text{Fe}_{2.7}\text{O}_{4-c}$

Neighbor	# of Neighbors	σ^2 ($\text{\AA}^2$)	Distance ($\text{\AA}$)	$\text{O}_h \text{Fe}_3\text{O}_4$ ($\text{\AA}$)	p
O	0.61(8)	0.001	1.720(8)	--	<0.001
O	5.42(8) ^b	0.0051(3)	2.019(5)	2.06	<0.001
Fe	4.07(4)	0.0039(9)	3.074(9)	2.97	<0.001
Tc	1.93(4) ^b	0.0039(9) ^c	3.074(9) ^c	2.97	0.004
Fe	6	0.0107(8)	3.509(9)	3.48	<0.001
O-Tc-O (MS)	6	0.014(2)	4.16(4)	4.12	0.001
Fe	12	0.014(2)	5.21(2)	5.14	0.003

- a) Fit range: $2 < k < 13.5$, $1 < R < 5.2$; 32.4 independent data. 13 parameters, $S_0^2 = 0.8$, $\Delta E_0 = 0(1)$ eV, $\chi^2 = 374$, $\chi^2_{\nu} = 21.5$, $R = 0.017$. Parameters without errors are fixed.
- b) Parameter is constrained to equal 6 minus the preceding parameter
- c) Parameter constrained to equal the preceding parameter

**Figure S48:** Tc K-edge EXAFS spectrum of Tc in $\text{Tc}_{0.1}\text{Ni}_{0.2}\text{Fe}_{2.7}\text{O}_{4-c}$

References

- (1) Lukens, W. W.; McKeown, D. A.; Buechele, A. C.; Muller, I. S.; Shuh, D. K.; Pegg, I. L. *Chem. Mater.* **2007**, *19*, 559.
- (2) Whitehead, D.; Malik, S. A. *Anal. Chem.* **1975**, *47*, 554.
- (3) Sugimoto, T.; Matijevic, E. *J. Colloid Interface Sci.* **1980**, *74*, 227.
- (4) Tang, Z. X.; Sorensen, C. M.; Klabunde, K. J.; Hadjipanayis, G. C. *J. Colloid Interface Sci.* **1991**, *146*, 38.
- (5) Koningsberger, D. C.; Prins, R. *X-Ray Absorption: Principles, Applications, Techniques of EXAFS, SEXAFS, and XANES*; John Wiley & Sons: New York, 1988.
- (6) Newville, M. J. *Synchrotron Rad.* **2001**, *8*, 322.
- (7) Ravel, B. *Phys. Scr.* **2005**, *T115*, 1007.
- (8) Ankudinov, A. L.; Ravel, B.; Rehr, J. J.; Conradson, S. D. *Phys. Rev. B* **1998**, *58*, 7565.
- (9) Bosi, F.; Halenius, U.; Skogby, H. *Am. Mineral.* **2009**, *94*, 181.
- (10) Downward, L.; Booth, C. H.; Lukens, W. W.; Bridges, F. *AIP Conference Proceedings* **2007**, *882*, 129.
- (11) Bluhm, H.; Andersson, K.; Araki, T.; Benzerara, K.; Brown, G. E.; Dynes, J. J.; Ghosal, S.; Gilles, M. K.; Hansen, H. C.; Hemminger, J. C.; Hitchcock, A. P.; Ketteler, G.; Kilcoyne, A. L. D.; Kneedler, E.; Lawrence, J. R.; Leppard, G. G.; Majzlan, J.; Mun, B. S.; Myneni, S. C. B.; Nilsson, A.; Ogasawara, H.; Ogletree, D. F.; Pecher, K.; Salmeron, M.; Shuh, D. K.; Tonner, B.; Tylliszczak, T.; Warwick, T.; Yoon, T. H. *J. Electron. Spectrosc. Relat. Phenom.* **2006**, *150*, 86.
- (12) Vanderlaan, G.; Kirkman, I. W. *Journal of Physics-Condensed Matter* **1992**, *4*, 4189.
- (13) Vanderlaan, G.; Thole, B. T. *Phys. Rev. B* **1991**, *43*, 13401.
- (14) Pattrick, R. A. D.; Van der Laan, G.; Henderson, C. M. B.; Kuiper, P.; Dudzik, E.; Vaughan, D. J. *Eur. J. Mineral.* **2002**, *14*, 1095.
- (15) Pearce, C. I.; Henderson, C. M. B.; Pattrick, R. A. D.; Van der Laan, G.; Vaughan, D. J. *Am. Mineral.* **2006**, *91*, 880.