

Supporting Information for:

Mn₂TeO₆: a distorted inverse trirutile structure.

Nami Matsubara^{1,2*}, Françoise Damay², Bénédicte Vertruyen³, Nicolas Barrier¹, Oleg I. Lebedev¹, Philippe Boullay¹, Erik Elkaïm⁴, Pascal Manuel⁵, Dmitry D. Khalyavin⁵, Christine Martin¹

¹*CRISMAT, 6 bvd Maréchal Juin, 14050 Caen Cedex, France*

²*Laboratoire Léon Brillouin, CEA-CNRS UMR12, Université Paris-Saclay, 91191 Gif sur Yvette Cedex, France*

³*GREENMAT, CESAM Research Unit, Université de Liège, Institut de Chimie B6, 13 Allée du 6 août, 4000 LIEGE, Belgium*

⁴*Synchrotron Soleil, Saint-Aubin BP 48, 91192 GIF-SUR-YVETTE CEDEX, France*

⁵*ISIS Facility, Rutherford Appleton Laboratory-CCLRC, Chilton, Didcot, Oxfordshire OX11 0QX, United Kingdom*

Figure S1 (color online): Description of the Σ_3 distortion components (polarization vectors) of all atoms in Mn₂TeO₆, using projections evidencing the two octahedra chains separately for simplicity. The direction and amplitude of the atomic displacements are schematized by arrows.

Figure S2 (color online): Projection along [001] of the RT structure of Mn₂TeO₆, deduced from the X-ray data refinement results, highlighting the oxygen atomic arrangement along *b* (blue ellipses) (for comparison with Figure 9d).




Figure S1




Figure S2